

PAPER NAME

ABS_190_Haryuni et al., 2023.docx

AUTHOR

Haryuni Haryuni

WORD COUNT

2576 Words

CHARACTER COUNT

13365 Characters

PAGE COUNT

5 Pages

FILE SIZE

52.0KB

SUBMISSION DATE

Jan 1, 2023 8:57 AM GMT+7

REPORT DATE

Jan 1, 2023 8:58 AM GMT+7

● 13% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 8% Internet database
- 11% Publications database
- Crossref database
- 8% Submitted Works database

● Excluded from Similarity Report

- Crossref Posted Content database
- Bibliographic material
- Quoted material
- Cited material

The effect of tobacco leaf biopesticides on the development bacterial leaf blight of rice

H Haryuni^{1*}, R Soelistijono¹, A F Aziez¹, I E Maryanti², S Supartini²

¹Faculty of Agriculture, Universitas Tunas Pembangunan Surakarta, Indonesia

²Faculty of Economic and Business, Universitas Tunas Pembangunan Surakarta, Indonesia

Corresponding author: haryuni@lecture.utp.ac.id

Abstract. The staple food of the Indonesian population is rice, the need for rice continues to increase following the increase in population, and one of the obstacles to increasing production is crop failure caused by rice leaf blight. Leaf blight is one of the important diseases which causes almost 70% yield loss. The purpose of this study was to determine the effect of the concentration and interval of application of tobacco leaf biopesticide on the development of leaf blight in Ciherang rice variety. This study was designed using a factorial complete randomized design, the first factor was the concentration of tobacco leaf biopesticide which consisted of 4 levels, namely 0 ml/l, 1 ml/l, 2 ml/l, and 3 ml/l. the second factor is the Giving Interval which consists of 3 levels. i. e. every 5 days, 10 days, and 15 days. Parameters of the development of rice bacterial leaf blight (BLB) that were observed were the number of grains per plant, percentage of attack and intensity of attack by leaf blight. The results showed that the highest number of grains per plant, the lowest attack percentage, and the lowest attack intensity were in the T3D1.

1. Introduction

The staple and main income of the Indonesian people come from agriculture, especially from rice commodities. Based on BPS data, the standard area of rice fields in Indonesia from 2021 to 2022 has increased from 10.41 million ha to 10.61 million ha, with a production of 54.42 million tons of GKG to 55.67 million tonnes of GKG with consumption in 2021 of 31.36 million tonnes in 2022 rising to 32.07 million tonnes, Indonesia's rice imports have increased from 2000 to 2021 of 407,741.4 tonnes [1].

Rice of the Ciherang variety is a superior variety, which is irrigated lowland rice, released in 2000, has a production range of 5-8 tons/ha, higher grain weight, and fluffier taste so that it is preferred by consumers, resistant to pests, and diseases [2], tolerant to floods and grown by more than 45% of farmers in Indonesia [3].

Rice is very susceptible to pathogens and affects the quality and quantity of yields, among several diseases that greatly impair rice yields [4] is bacterial leaf blight (BLB) caused by *Xanthomonas oryzae* pv. *oryzae* strains III and IV are the most important rice diseases in the world [5], another term for crackle disease in rice [4], seed dispersal, fast spread, broad spectrum causing huge losses reaching around 30 - 70%, crop failure or loss [6] symptoms that cause grain rot [7], gray to white lesions along the leaf veins or vascular disease, vascular infection [4] through seeds, natural holes in plants, wounds develop in xylem vessels [8]. The population is difficult to control, has many pathotypes, and is dynamic so that it damages plant resistance [9].

Control of BLB is carried out at the beginning of growth before the formation of tillers, stopping it will result in a decrease in production, efforts to control organic farming are environmentally friendly by utilizing plants that have the potential as antipathogens [10]. Some plant species have toxic properties, can control plant pathogens, are easy to decompose and are safe for a long time, however, it takes longer to kill pathogens than using chemicals [11]. Types of plants that can be used as vegetable pesticides include lemongrass, soursop, tobacco, and so on, these plants have bioactive compounds that act as repellents, deterrents and attracting pests and diseases [12]. Several researchers tested tobacco leaf biopesticide for controlling plant pests *Heliothis virescens* [13] *Myzus persicae*. Sulz [14], *Hypothenemus hampei* [15]. Tests for BLB on the Ciherang rice variety have never been carried out. The purpose of this study was to determine the effect of the concentration and interval of application of tobacco leaf biopesticide on the development of leaf blight in Ciherang rice variety.

1. Material and methods

The study was conducted from May to September 2020. The experiment in Kwangsang village. Jumapolo, Karanganyar, Central of Java. Laboratory analysis was carried out at the Plant Agroteknology Laboratory, Faculty of Agriculture, Universitas Tunas Pembangunan Surakarta (UTP).

The tobacco leaf biopesticide used Sentosa method [16], produced from research group between Universitas Indonesia and PT. Zekindo, The Rice Seed (*Oryza sativa*) used in the Ciherang variety by the Food Crop Seed Certification Supervision Agency. The Ciherang variety rice seeds were soaked for 12 hours then drained then waited for 2 days until shoots appeared, after 14 days after planting then the seeds were sown. A week before planting, tillage is carried out, consisting of two stages, namely turning the soil over and leveling the land [17].

This study was designed using a factorial complete randomized design, the first factor was the concentration of tobacco leaf biopesticide (T) which consisted of 4 levels, namely 0 ml/l, 1 ml/l, 2 ml/l, and 3 ml/l. the second factor is the giving Interval (D) which consists of 3 levels. i. e. every 5 days, 10 days, and 15 days. Data were analyzed using analysis of variance (ANOVA) and if they were significantly different, they were further tested with DMRT with a 95% confidence level [18].

3. Result and Discussion

Observations of the effect of the average concentration and interval application of tobacco leaf biopesticide on Ciherang variety rice were analyzed by Duncan multiple range test (DMRT) at 5% level presented in Table 1.

Table 1. The effect of the concentration of tobacco leaf biopesticides on the number of grains per plant, tillers per clump, attack percentage, and attack intensity.

Treatment	Parameters			
	Number of grains per plant (grain)	Tillers per clump (stem)	Attack percentage (%)	Attack intensity (%)
Concentration of biopesticide				
T0	627.04 ^a	11.16	16.53 ^b	16.88 ^b
T1	780.92 ^b	10.63	10.36 ^a	10.31 ^a
T2	888.02 ^b	11.01	9.08 ^a	9.01 ^a
T3	976.44 ^d	10.66	7.66 ^a	8.31 ^a

Note. Numbers followed by the same letter show no significant difference in the Duncan multiple range test (DMRT) 5% level.

Description: T0= tobacco leaf biopesticide 0 ml /L, T1= tobacco leaf biopesticide 1ml /L, T2= tobacco leaf biopesticide 2 ml /L, T3= tobacco leaf biopesticide 3 ml /L.

Table 1 on the observation of the number of grains of rice per plant showed a significant difference between T0 and T1, T2, T3, the highest number of grains per plant in the T3 treatment was significantly different from T0, T1 and T2. the concentration of tobacco leaf biopesticide affects the number of grains per plant, info for help [17] tobacco leaf biopesticide contains 33.08% nicotine. Nicotine is produced from tobacco roots as a secondary metabolite which is excreted through root exudation in the rhizosphere area thereby increasing the uptake of nitrogen, phosphorus, potassium [19] zinc, and iron [20]. In addition to the levels of these elements, tobacco plants also produce gibberellin hormones found in rice plants. Gibberellin plays a role in synthesizing the enzyme α -amylase which stimulates starch hydrolysis and increases cell elongation which in turn affects the number of grains per plant [21].

Observation of tillers per clump did not differ significantly between treatments indicating that the concentration of tobacco leaf biopesticide had not had an effect, possibly the microclimate that caused it was not significantly different [22], research on the concentration of tobacco leaf biopesticide still needs to be continued to obtain more tangible results.

The percentage and intensity of BLB attacks (table 1) showed that T0 was significantly different from T1, T2, and T3. The highest attack without tobacco leaf biopesticides (T0). Increasing the concentration of tobacco leaf biopesticides reduced the percentage of BLB attacks. Tobacco leaf biopesticide contains methanol compounds which act as antibacterial [23]. It is supported by the research of [24] which states that methanol content with a concentration of 25 mg/ml acts as an antimicrobial against 6 tested bacterial isolates.

Table 2. The effect of interval of tobacco leaf biopesticide on the number of grains per plant, tiller per clump, attack percentage, and attack intensity.

Treatment	Parameters			
	Number of rice grains per plant (grain)	Tillers per clump (stem)	Attack percentage (%)	Attack intensity (%)
Interval of tobacco leaf biopesticide				
D1	889.68 ^b	11.00	11.36 ^b	10.47 ^a
D2	799.41 ^b	10.53	10.97 ^b	11.27 ^b
D3	765.23 ^a	11.06	11.36 ^b	11.73 ^b

Note: Numbers followed by the same letter show no significant difference in the Duncan multiple range test (DMRT) 5% level.

Description: D1= every 5 days, D2= every 10 days, D3= every 15 days.

Table 2 shows that the intervals for administering tobacco leaf biopesticides D1 and D2 were significantly different from D3. At intervals of 15 days the number of grains decreased because the gibberellin hormone produced by tobacco leaf biopesticides was not sufficient for its needs so longer intervals reduced the number of grains while observing tillers per clump at intervals did not give a significant effect.

Observation of attack percentage and intensity of BLB attack on Ciherang variety showed a significant difference 5 days interval (D1) was lower than D2 and D3. This is because the content of methanol and phenolic compounds is sufficient to suppress the development of BLB [25].

Table 3 shows that the highest number of seeds planted was in the T3D1 treatment, the highest tillers per clump were T1D3, while the lowest attack percentage and attack intensity were T3D1. Tobacco leaf biopesticides stimulate the formation of gibberellin which plays a role in the process of plant photosynthesis and then facilitates rapid cell division [26]. Increasing the concentration and increasing the time interval every 15 days of tobacco leaf biopesticides were not significantly different, the best conditions were achieved at high concentrations and intervals every 5 days were able to reduce the percentage and intensity of attacks.

Table 3. The effect interaction of the concentration and interval of tobacco leaf biopesticides on the number of grains per plant, tillers per clump, attack percentage, and attack intensity.

Treatment	Parameters			
	Number of grains per plant (grain)	Tillers per clump (stem)	Attack percentage (%)	Attack intensity (%)
Interaction of the concentration and interval of tobacco leaf biopesticides				
T0D1	647.27 ^a	10.63	17.17 ⁱ	17.07 ^e
T1D1	911.37 ^{cd}	11.30	9.50 ^{def}	9.47 ^{bc}
T2D1	994.50 ^{de}	11.20	8.30 ^{bc}	8.40 ^b
T3D1	1,005.60 ^e	10.87	6.60 ^a	6.93 ^a
T0D2	591.73 ^a	10.87	16.80 ⁱ	17.07 ^e
T1D2	767.23 ^{bc}	10.97	10.30 ^{fg}	10.23 ^{cd}
T2D2	852.60 ^{bc}	9.93	9.07 ^{cde}	9.40 ^{bc}
T3D2	986.07 ^{de}	10.63	15.63 ^h	8.40 ^b
T0D3	642.13 ^a	10.60	11.27 ^g	16.50 ^e
T1D3	664.17 ^a	11.87	11.27 ^h	11.23 ^d
T2D3	816.97 ^b	10.67	9.87 ^{ef}	10.13 ^{cd}
T3D3	937.67 ^{cde}	11.20	8.67 ^{bcd}	9.08 ^{bc}

Note: Numbers followed by the same letter show no significant difference in the Duncan multiple range test (DMRT) 5% level.

Description: T0= tobacco leaf biopesticide 0 ml /L, T1= tobacco leaf biopesticide 1 ml /L, T2= tobacco leaf biopesticide 2 ml /L, T3= tobacco leaf biopesticide 3 ml /L. D1= every 5 days, D2= every 10 days, D3= every 15 days.

Tobacco leaf biopesticide has antimicrobial activity as an antibacterial [20,27] antifungal [20,25]. The highest biopesticide content in tobacco leaves is nicotine and several N-containing compounds or groups such as 3-(1-methyl-2-pyrrolidiny)-pyridine which play a role in increasing nutrient absorption in vanilla [19] and corn plants [20]. Nitrogen for plants is an important component of protein, chlorophyll and genetic material so that it has a direct effect on growth, reproduction and inhibits the formation of pathogenic chitin *Fusarium oxysporum* f.sp vanilla [20].

4. Conclusion

Based on the results of data analysis using the Anova test and Duncan's 5% follow-up test, the effect of the concentration and interval of tobacco leaf biopesticide on the Ciherang rice variety had no significant effect on tillers per clumps but had a significant effect on the number of grains per plant, the percentage of BLB attacks and the intensity of BLB attacks. The treatment that showed the highest number of grains (1,005.60 grains, the lowest percentage of attack was 8.60%, and the lowest intensity of attack was 6.93% in the combination treatment of 3ml/l of tobacco leaf biopesticide concentration and spraying interval every 5 days (T3D1).

References

- [1] BPS, 2022 <https://www.bps.go.id/pressrelease/2022/10/17/1910/pada-2022> accessed December 15, 2022.

- [2] Anonymous. 2019. Agricultural Research and Development Agency. Agricultural Research and Development Agency. <http://www.litbang.pertanian.go.id/varietas/130/> accessed December 25, 2022.
- [3] Nugraha Y, Hidayatun N, Trisnaningsih, Yuliani D, Ardiyanti S and Kadir T S. 2017. *Indonesian Journal of Agricultural Science* **18**.1: 7–1
- [4] Sharma Y, Dua D and Nagar A, Srivastava NS. 2016. *Int J Pharm Sci Res* **7**.3: 1156–67.
- [5] Zhang H and Wang, S. 2013. *Current Opinion in Plant Biology*, **16**.2:188–95.
- [6] Weny I Safni, Lisnawita and Lubis K .2019. IOP Conf. Ser.: Earth Environ. Sci. **260** 012118
- [7] Joko T, Kristantini, Sumarno, and Andriyanto R. 2019. *Jurnal Perlindungan Tanaman Indonesia*. **23**.2: 205–210
- [8] Zhao Y, Qian, Fan G, Yin J, Zhou Y, Liu Y, Shen C, Hu Q and Liu F. 2012. *Phytopathology* 102: 252–9
- [9] Sudir, Nuryanto B and Kadir T S. 2012. *Iptek Tanaman Pangan* 7: 79–87.
- [10] Sati SC and Joshi S. 2011. *Research Journal of Medicinal Plant* **5**.4: 377–91.
- [11] Yellareddygarri S KR, Vijay-Krishna-Kumar K, Sudini H, Reddy E C S, Sayyed R Z, Naga-Madhuri K V, Suresh K, Ravindra-Reddy B, Hemalatha S, Al-Turki A, El-Meleigi M A, Han Y, Sruthi V S and Reddy M S.. Chapter 42. <https://www.researchgate.net/publication/274068381> .
[aaccessed 12/27/2022](https://www.researchgate.net/publication/274068381)
- [12] Wuryantini S, Harwanto and R A Yudistira. 2020 IOP Conf. Ser: Earth Environ. Sci. **457** 012039
- [13] Viteri D M, Sarmiento L, Linares A L, and Cabrera, I. 2019. *Crop Protection*, **122**, 175–9. <https://doi.org/https://doi.org/10.1016/j.cropro.2019.05.008>
- [14] Khan I A, Hussain S, Akbar R, Saeed M, Farid A, Ali I, Alam M, Shah B. 2015. *Journal of Entomology and Zoology Studies*. **4**: 371–3.
- [15] H. Haryuni, T. S. K. Dewi, E. Dewi, S. F. Rahman, and M. Gozan. 2019. *International Journal of Technology*, **10** 1:159–66.
- [16] Sentosa Y, H N Andjani, K Yati, M Jufri, Haryuni and M Gozan 2019 Determination of LC50 Value of *Nicotiana tabacum* L. Extract Against *Tenebrio molitor* and *Zophobas morio* Larvae in AIP Conference Proceedings, 2193 1: 030021
- [17] Kartahadimaja J, Wentasari R, and Sesanti R N 2010 *Agrovigor* 3 131–13
- [18] Gomez, K.A. and Gomez, A.A. 1995. *Prosedur Statistika untuk Penelitian Pertanian Edisi Kedua* (Endang Sjamsuddin & Justika S. Bahrsjah. Terjemahan). Jakarta: UI Press. (in Indonesia)
- [19] Haryuni, H., Harahap, A. F. P., Supartini, Priyatmojo, A., & Gozan, M.. *International Journal of Agronomy*, 1–6.
- [20] Lisuma, J. B. ; Mbega, E. R. ; Ndakidemi, P. A.. 2019. *International Journal of Agriculture and Biology* **22** 1: 1–12
- [21] Utama R C and Sugiyatna. 2016. *Bul Agrohorti*, **4** 1: 56–62.
- [22] Sumardi D, Bahariawan M, Maulani R R, Suhandono S, Novia C, Harahap A F E and Gozan. 2021. IOP Conf. Ser.: Earth Environ. Sci. 940 012026
- [23] Singh K P, Daboriya V, Kumar S, Singh S. *Rom. J. Biol. – Plant Biol.* **55** 2: 135–142
- [24] Akinpelu DA, Obuotor EM.2000. *Fitoterapia*.**71** 2:199–200
- [25] Duan CB, Du Y, Hou X, et al. 2016. *Molecules*. **21**12: 1743
- [26] Mona M A 2013 *Int. J. Plant Physiol. Biochem.* 5: 42–49
- [27] Ameya G, Manilal A, Merdekios B. 2017. *Open Microbiol J.* 11: 352–9

13% Overall Similarity

Top sources found in the following databases:

- 8% Internet database
- 11% Publications database
- Crossref database
- 8% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	H Haryuni, I Y Atmaja, T Supriyadi, S Priyadi, M Rumaningsih. "Effect of..."	Crossref	4%
2	S J Rachmawatie, E Purwanto, A T Sakya, W S Dewi. "Growth and conte..."	Crossref	1%
3	Higher Education Commission Pakistan on 2018-06-25	Submitted works	<1%
4	Robbani Hanida, Sih Dewi Widyatmani, Prasojo Haryuni, Supriyadi. "Im..."	Crossref	<1%
5	Wathome N. Mbatha, Nicholas K. Korir, Joseph P. Gweyi-Onyango. "Infl..."	Crossref	<1%
6	biodiversitas.mipa.uns.ac.id	Internet	<1%
7	D Sumardi, M Bahariawan, R R Maulani, S Suhandono, C Novia, A F P H...	Crossref	<1%
8	Higher Education Commission Pakistan on 2015-10-17	Submitted works	<1%
9	journal.unnes.ac.id	Internet	<1%

10	worldwidescience.org Internet	<1%
11	repository.ipb.ac.id Internet	<1%
12	Achmad Fatchul Aziez. "Maximizing the Yield of Black Cute Rice using ... Crossref	<1%
13	jurnal.upnyk.ac.id Internet	<1%
14	ncbi.nlm.nih.gov Internet	<1%
15	hindawi.com Internet	<1%
16	hdl.handle.net Internet	<1%