

## COVERING LETTER

Manuscript has main author and co-authors. Author names should not contain academic title or rank. Indicate the corresponding author clearly for handling all stages of pre-publication and post-publication. Consist of full name author and co-authors. **Corresponding author** is a person who is willing to handle correspondence at all stages of refereeing and publication, also post publication.

### I. First author\*:

1. Name : Haryuni Haryuni
2. Affiliation : Fakultas Pertanian Universitas Tunas Pembangunan Surakarta
3. E-mail : haryuni@lecture.utp.ac.id
4. Orcid ID : <http://orcid.org/0000-0002-2574-5506>
5. Phone number : 0812-297-7555

### II. Second author:

1. Name : Muhamad Amin
2. Affiliation : Mahasiswa Fakultas Pertanian Universitas Tunas Pembangunan Surakarta
3. E-mail : [aminutp2015@gmail.com](mailto:aminutp2015@gmail.com)
4. Orcid ID : -
5. Phone number : 0813-3228-3690

### III. Third author:

1. Name : Endang Suprpti
2. Affiliation : Fakultas Pertanian Universitas Tunas Pembangunan Surakarta
3. E-mail : [endang.suprpti@lecture.utp.ac.id](mailto:endang.suprpti@lecture.utp.ac.id)
4. Orcid ID : -
5. Phone number : 0813-2944-5408

### IV. Fourth author:

1. Name : Tyas Soemarah Kurnia Dewi
2. Affiliation : Fakultas pertanian Universitas Tunas Pembangunan Surakarta
3. E-mail : [tyas.soemarah@lecture.utp.ac.id](mailto:tyas.soemarah@lecture.utp.ac.id)
4. Orcid ID : -
5. Phone number : 0815-4841-9240

### V. Fifth author

1. Name : Eko Hartoyo
2. Affiliation : Fakultas pertanian Universitas Tunas Pembangunan Surakarta
3. E-mail : [hartoyoeko56@gmail.com](mailto:hartoyoeko56@gmail.com)
4. Orcid ID : <https://orcid.org/0000-0003-2358-8452>
5. Phone number : 0813-2869-2250

**VI. Sixth author:**

1. Name : Mrihrahayu Rumaningsih
2. Affiliation : Fakultas Ekonomi dan Bisnis Universitas Tunas Pembangunan Surakarta
3. E-mail : [mrihrahayu.rumaningsih@lecture.utp.ac.id](mailto:mrihrahayu.rumaningsih@lecture.utp.ac.id)
4. Orcid ID : -
5. Phone number : 0813-2901-9012

**VII. Seventh author**

1. Name : Supartini Supartini
2. Affiliation : Fakultas Ekonomi dan Bisnis Universitas Tunas Pembangunan Surakarta
3. E-mail : [supartini@lecture.utp.ac.id](mailto:supartini@lecture.utp.ac.id)
4. Orcid ID : -
5. Phone number : 0813-2907-2731

**VIII. Eight author**

1. Name : Istinganah Eni Maryanti
2. Affiliation : Fakultas Ekonomi dan Bisnis Universitas Tunas Pembangunan Surakarta
3. E-mail : [istinganah.maryanti@lecture.utp.ac.id](mailto:istinganah.maryanti@lecture.utp.ac.id)
4. Orcid ID : <https://orcid.org/0000-0002-2995-0435>
1. Phone number : 0812-2975-6690

# PROLINE REDUCTION AND INCREASING GROWTH OF VANILLA PLANTS INDUCED BY BNR FUNGUS WITH DOSE OF GOAT MANURE AND HUSK CHARCOAL

Haryuni Haryuni<sup>1\*</sup>, Muhammad Amin<sup>1</sup>, Endang Suprapti<sup>1</sup>, Tyas Sumarah Kurnia Dewi<sup>1</sup>, Eko Hartoyo<sup>2</sup>, Mrihrahayu Rumaningsih,<sup>3</sup> Supartini Supartini<sup>3</sup>, Istinganah Eni Maryani<sup>3</sup>,

<sup>1</sup> Department of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta, Central of Java, Indonesia

<sup>2</sup> Department of Agribisnis, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta, Central of Java, Indonesia

<sup>3</sup> Department of Management, Faculty of Economy and Bisnis Universitas Tunas Pembangunan, Surakarta. Central of Java, Indonesia

\*Corresponding author: haryuni@lecture,utp.ac.id

## Abstract

Goat manure and husk charcoal are agricultural wastes that have not been used optimally, contain soil microorganisms that are effective in converting nutrients into more perfect forms, play a role in improving soil physical properties and increase the ability of soil to bind nutrients. The purpose of this study is to determine the effect of goat manure and husk charcoal to proline reduction and increase growth of vanilla plant induced by Binucleate Rhizoctonia (BNR). The study was designed with a factorial completely randomized block design. The first factor is the dose of manure (0,5,10,15) g/plant, the second factor is the dose of husk charcoal (0,5, 10, 15) g/plant. Data were analyzed by analysis of variance ANOVA with Duncan's Multiple Distance Test (DMRT) with a significant difference of 5%. The effect of doses of goat manure and husk charcoal induced by BNR can reduce proline contents and increase growth of vanilla plant. The proline content reduce to affects the resistance of vanilla plants to drought.

**Keywords:** binucleate-Rizoctonia, husk, manure, vanilla

## INTRODUCTION

Vanilla is one of the plantation crops which is the country's foreign exchange reserves from exports. Indonesian vanilla comes from Mexico, the first use of vanilla was done by the Aztecs as a preserved chocolate fragrance that enhances the taste. Vanillin ( $C_8H_8O_3$ ) content which causes vanilla (Nurcholis, 2017) is a distinctive fragrance used in the food, beverage, cosmetic, and pharmaceutical industries. Vanilla plants are characterized by growth that propagates up to 30 meters, epiphytes and vanilla fruit producing pods (Frenkel & Belanger, 2018). The initial growth of vanilla requires a medium that has the ability to

hold water, loose soil structure, smooth aeration and drainage, does not experience drought and contains sufficient nutrients needed for growth (Fileccia et al., 2014)

Fertilization treatments, especially organic fertilizers, accelerate soil decomposition so that soil microorganisms can help provide nutrients needed by plants. Organic fertilizer derived from goat manure has the advantage of increasing soil humus content, stimulating soil microorganisms, and improving soil structure so as to increase the availability of nutrients for plants (Singh, 2012); (Assefa & Tadesse, 2019); (Pratomo & Prasetyo, 2018). Application of organic matter derived from goat manure enriches microorganisms, produces nutrients (macro and micro), increases soil aggregates so as to increase water infiltration and acts as a soil enhancer. Goat manure has a long-term effect on Potassium content in goat manure is 2.88%, other nutrients nitrogen is 2.77%, phosphorus is 1.78% (Handajaningih et al., 2018).

In addition to goat manure, soil organic matter can also be added by giving husk charcoal. Treatment with husk charcoal increased growth and reduced the percentage of seedling mortality caused by the fungus *Fusarium oxysporum* f. sp. *Vanilla* (Haryuni et al., 2018). The results of the decomposition of organic matter from goat manure and husk charcoal (N, P, K, Ca, S and Mg) which were eliminated directly were able to increase the pH and increase the ability of the soil as a cation buffer because the decomposition of organic matter produced a complex compound (humus) (Fileccia et al., 2014). Organic fertilizer derived from goat manure has the advantage of increasing soil humus content, stimulating soil microorganisms, and improving soil structure so as to increase the availability of nutrients for plants. Continuous use of land without being followed by nutrient supply, it is difficult to obtain maximum production considering the very limited capacity of the soil, the provision of goat manure with an appropriate dose is expected to help the availability of soil nutrients, goat manure and husk charcoal ash serve as biofertilizers (Harahap & Hidayat, 2021).

Plant growth is influenced by the availability of nutrients that can be utilized by roots and available to plants, this role can be carried out by Binucleate Rhizoctonia (BNR). The peloton that is in the root tissue plays a role in storing nutrients that are absorbed by external hyphae and sent into plant cells (Haryuni, et al., 2020). The characteristic preserved mycorrhizal fungi associated with orchids is the formation of a symbiotic structure called the peloton, which is shaped like a coil of hyphae in which metabolic exchange between symbionts occurs in both protochrome and mycorrhizal roots (Valadares et al., 2014). The

material in the peloton that is in the network is utilized by plants during the metabolic process. In conditions of water shortage, the material contributes to the availability of water for plants (Begum et al., 2019). BNR-mediated mechanisms regulate tolerance and promote plant improvement in drought stress conditions (Boutasknit et al., 2020).

Proline is a compatible osmolyte commonly found in plants experiencing drought stress. Proline accumulation is a typical physiological response to drought stress in some species. (Kumar et al., 2021). Proline is an indicator of a plant's ability to maintain turgor through a decrease in osmotic potential so that an increase in proline content indicates an increase in plant tolerance to water shortages, which results in plant growth and the quantity and quality of crop yields. Proline affects the air potential, opening and closing of leaf stomata further down regulates photosynthesis and CO<sub>2</sub> deficiency causes plant stress (Chun et al., 2018); (Osakabe et al., 2014). There is not enough information about the application of goat manure and husk charcoal of vanilla induce by BNR and their effect to proline contents and plant growth, this research can provide this information.

## **METHODS**

Preparation of goat manure, the goat manure used is derived from wind-dried goat manure until the moisture content reaches 60%, put in a wooden box then added with molasses/Javanese sugar and coconut water in a ratio (10:1:3) mixed evenly and stored closed, every 7 days once inverted and stirred for 5 times. If the fertilizer has changed color and comes off when held by hand, the fertilizer is ready to use (Pratomo & Prasetyo, 2018).

Preparation of husk charcoal, the husk charcoal is cleaned of other impurities and collected together and then a chimney/tube is installed in the middle which is made of perforated wire. wire is shaped like a tube as a means of combustion. This means that the flames will stick out. For burning, it is better to use a base made of zinc plate placed on a flat, hard and heat-resistant floor. The hole in the chimney is filled with charcoal and doused with kerosene, added with paper, so that a fire arises, when it is black the unburned part is piled up and moved near the chimney. After everything is burned, the husks turn black, allow to cool and are ready to be used for planting media mixtures (Surdianto et al., 2015).

Preparation of Binucleat Rhizoctonia (BNR), the BNR used is the same as in the (Haryuni et al., 2015 b) procedure, namely the corn is washed, drained and then steamed for 45 minutes until half, it is half cooked, cooled, then sprayed with water that has been mixed

with chlorine, broken corn and then put into a heat-resistant plastic bag with a size of 10 x 20 cm until it reaches 1/3 part . After being put in a small plastic bag, stored in a large plastic bag to be sterilized for 60 minutes by steaming, cooling at room temperature. Pure isolates of BNR were propagated on PDA media, inoculated to 100g sterile corn medium and stored at room temperature for 10-15 days in a separate room not used for other activities. After the corn medium was filled with BNR growth, it was inoculated into the vanilla growing medium. Each vanilla plant was inoculated with 10 g of BNR before planting.

The research was conducted at Rejosari village, Jambu district, Semarang Regency, Central of Java from March to September 2020 the altitude is around 800-900 m above sea level, soil type andosol soil. Vanilla plants that are 2 years old in the field, sprinkled with BNR 10 g/plant around the plant roots by covering the surface soil, after 2 weeks of giving BNR goat manure is given at a dose of 0.5,10,15 g/plant as (P0 , P1, P2, P3) every 2 weeks up to 4 times. The second factor was the dose of husk charcoal 0,5,10,15 g/plant as (S0, S1, S2, S3) added every 2 weeks up to 4 times. each treatment was repeated 5 times used samples.

Growth Observation of vanilla plant was carried out 4 weeks after the first treatment of goat manure and husk charcoal including, plant height (measured from the base of the stem to the top of the stem), number of leaves (counted for all the leaves that grew for each sample plant), leaf fresh weight (taken 10 leaves). leaf starting with 5 leaves from the stem surface and then weighed), leaf dry weight all leaves were observed on leaf fresh weight by drying the leaves and continued in an oven at 105<sup>o</sup>C for 2 h and continuously at 80<sup>o</sup>C until the weight was constant (Huang et al., 2019).

Measurement of Proline contents, analysis of proline contents in leaf of vanilla was determined following (Bates et al., 1973). Extract: 20 to 50 times diluted fresh weight of leaf (w/v), typically in a 70:30 ethanol:water mixture (v/v). Standards: proline solutions ranging from 0.04 to 1 mM, in the same medium as the one used for the extraction. Reaction mix: ninhydrin 1% (w/v) in acetic acid 60% (v/v), ethanol 20% (v/v). Protect from light. Proline analysis on Laboratory of fisiology at Universitas Gadjah Mada Yogyakarta.

### **Statistical analysis**

Data from analysis of proline and vanilla plant growth (height, fresh weight and dry weight of plants, analyzed by variance (Anova). If significantly different, Duncan's Multiple

Range Test (DMRT) test at 5% level was carried out, to distinguish between treatments. To find out the direct and indirect relationship, regression and correlation analysis were carried out.

## RESULT

The average effect of goat manure on proline content, height, fresh weight and dry weight of vanilla plants induced by BNR fungus analyze the duncan multiple range test (DMRT) 5% level in activities in experiment and control is presented in Table 1.

Table 1. Effect of goat manure to proline content, height, fresh weight and dry weight of vanilla plants induced by BNR fungus.

Treatment	Observation of parameters			
	Proline content ( $\mu\text{mol g}^{-1}$ )	height of plant (cm)	fresh weight of plant (g)	dry weight of plant (g)
P0	2.12 b	50.17 a	6.05 b	1.27 c
P1	2.06 b	58.71 ab	6.14 b	1.31 b
P2	1.91 ab	64.88 bc	6.19 ab	1.34 a
P3	1.77 a	71.08 c	6.37 a	1.36 a

Note: Numbers followed by the same letter show no significant difference in the duncan multiple range test (DMRT) 5% level.

Goat manure has a significant effect on proline content, height, fresh weight, and dry weight of vanilla plant (Tabel 1). It caused by goat manure contains macro and micro nutrients rich in phosphorus, nitrogen and potassium which are needed for plant growth and development. Increasing the dose of goat manure decreased and significantly different on plant proline content (Yenni et al., 2021), are (P0) 2.21  $\mu\text{mol g}^{-1}$ , (P3) 1.77  $\mu\text{mol g}^{-1}$ . But it tends to increase and significantly different on height of vanilla plant (P0) 50.17cm, (P3) 71.08cm. the fresh weight of plant (P0) was 6.05 g, (P3) 6.37g, and the dry weight of plant (P0) was 1.27 g, (P3) 1.36 g. There are caused by BNR fungi on plants acts as mycorrhizae capable of forming internal and external hyphae on roots and peloton in plant tissues (Haryuni. et al., 2020). Table 1 shows that increasing the dose of goat manure fertilizer decreased plant proline content and increased plant height, plant fresh weight, plant dry weight.

Glutamate is an important amino acid and acts as a precursor for biosynthesis - aminobutyric acid, arginine, glutamine, and proline in plants and other eukaryotes, proline as an osmoprotectant in stress tolerance. Proline participates in biosynthesis primary

metabolism, but also has a special function, as a metabolite during growth and development, Proline not only accumulates during abiotic stress, but also in various plant tissues under non-stress condition too (Kavi Kishor et al., 2015). Proline is an indicator of plant stress against drought tolerance of peanut plants (Furlan et al., 2020) with the addition of goat manure to help soil microorganisms more readily utilized by plants and then stored in plant roots and stems in pelotons produced from BNR fungus inoculation. Fungal hyphae penetrate cortical cells, branch and form peloton (cells in this phase are called trophocytes), as an anatomical feature of orchid mycorrhizae

This is the same as research carried out by (Zhang et al., 2016) that BNR application as mycorrhizal can increase total root long on the black grasshopper plan. BNR fungus forms external and internal hyphae on roots and forms peloton in plant transport bundles, microorganisms contained in goat manure are able to improve soil structure so that water in the soil is more easily absorbed by plant roots. Lack of water of vanilla plant tissues is indicated by high contents of proline (Haryuni et al., 2020).

Accumulation of free proline as a form of general response of a plant to stress factors in drought, salinity, cold air and pathogen. Height, fresh weight and dry weight are indicators of plant growth and development of the plant, groundwater that enters through the roots is stored in plant tissue, used in metabolic processes and used for plant growth and development. Lack of water in plants will affect physiological processes and even cause plants to die (Pardosi, 2021).

The average effect of husk charcoal to proline content, height, fresh weight and dry weight of vanilla plants induced by BNR fungus analyze the duncan multiple range test (DMRT) 5% level in activities in experiment and control is presented in Table 2.

Table 2. Effect of husk charcoal dose to proline content, height, fresh weight and dry weight on vanilla plants induced by BNR fungus

Treatment	Observation of parameters			
	Proline content ( $\mu\text{mol g}^{-1}$ )	height of plant (cm)	fresh weight of plant (g)	dry weight of plant (g)
S0	3.59 d	25.04 a	5.41 d	1.13 d
S1	2.14 c	37.83 b	6.13 c	1.28 c
S2	1.27 b	73.21 c	6.50 b	1.40 b
S3	0.85 a	108.75 d	6.70 a	1.47 a

Note: Numbers followed by the same letter show no significant difference in the DMRT test at 5% level.



Table 2 shows the significant effect between the application treatment of husk charcoal and without husk charcoal to proline content, plant height, weight, and dry weight of plant. Proline contents were significantly different and decreased with the application of husk charcoal at a dose of 15 g/plant (S3) by  $0.85 \mu\text{mol g}^{-1}$  all (S0)  $3.59 \mu\text{mol g}^{-1}$  but tended to increase and significantly different in plant height from 25.04 cm (S0) to 108.75 (S3). the weight of the wet stove from 5.41 g (S0) to 6.70 g (S3) and the dry weight of 1.13 g (S0) to 1.47 g (S0).

The treatment of increasing doses of husk charcoal charcoal, proline has increased, in accordance with the opinion of (Hendrati et al., 2016), proline will increase in tolerant plants not in sensitive plants, so that proline can be used as an indicator of drought tolerant from *Acacia auriculiformis* Cunn., *Tectona grandis* L., *Alstonia spectabilis* Br., and *Cedrela odorata* L. that are susceptible to drought. resistant to drought does not produce proline. Supported by the opinion of (Ichsan et al., 2016), The role of husk charcoal as soil organic matter stimulates soil granulation, cohesion and plasticity, improves soil structure and increases soil resistance to water and increases activity and number of soil microorganisms. The increase in soil organisms was supported by the induction of the BNR which was inoculated at the time of *Shorea robusta* seedlings (Tapwal et al., 2015).

The increase in the fresh weight of plant was indicated by an increase in the dose of husk charcoal charcoal, the fresh weight was related to the photosynthate of the plant that was stored in the tissues and the absorption of water through the roots, the availability of nutrients and water in the soil affected the growth and fresh weight. Dry weight is an indicator of photosynthetic accumulation of plants stored in soybean plant tissue used for growth and development (Widiastuti & Latifah, 2016). However, used leaf dry weight as the representative of leaf biomass when studying the scaling relationship between leaf biomass and area. The scaling of leaf mass and leaf area could directly influence the amount of light-capturing surface area, which gives an investment of dry biomass (Huang et al., 2019).

Determine leaf biomass related to leaf moisture content, by measuring leaf dry weight and leaf fresh weight so that it can determine water loss, to make it easier to determine gas exchange ( $\text{CO}_2$ ,  $\text{O}_2$ , and  $\text{H}_2\text{O}$ ) leaf dry weight must be constant. (Hughes et al., 1970) reported that leaf area of several dicot genotypes had a linear relationship with absolute leaf water content (Lin et al., 2018) described the comparison between fresh leaf weight and leaf area of 11 bamboo species.

Table 3. The interaction effect of goat manure and husk charcoal dose on proline content, height, fresh weight and dry weight to vanilla plants induced by BNR fungus

Treatment	Observation of parameters			
	Proline content ( $\mu\text{mol g}^{-1}$ )	height of plant (cm)	fresh weight of plant (g)	dry weight of plant (g)
P0S0	3.94 j	22.33 a	5.20 f	1.05 h
P1S0	3.45 hi	25.00 a	5.23 f	1.10 h
P2S0	3.65 i	25.00 a	5.33 f	1.16 g
P3S0	3.33 h	27.83 a	5.90 e	1.20 fg
P0S1	2.35 g	30.00 ab	6.00 ed	1.23 ef
P1S1	2.34 g	33.67 ab	6.13 cde	1.26 de
P2S1	2.37 g	40.33 bc	6.17 cde	1.30 cd
P3S1	1.51 f	47.33 cd	6.23 bcde	1.32 cd
P0S2	1.09 bc	54.00 d	6.40 abcd	1.34 c
P1S2	1.21 cde	73.50 e	6.50 abc	1.40 b
P2S2	1.42 ef	80.33 e	6.53 abc	1.42 b
P3S2	1.35 def	85.00 ef	6.60 ab	1.43 ab
P0S3	1.11 bcd	94.33 fg	6.63 ab	1.47 a
P1S3	0.64 a	102.67 g	6.70 a	1.47 a
P2S3	0.77 a	113.83 h	6.73 a	1.47 a
P3S3	0.89 ab	124.17 h	6.77 a	1.47 a

Note: Numbers followed by the same letter show no significant difference in the DMRT test at 5% level.

Table 3 shows the interaction effect of goat manure and husk charcoal dosage charcoal induced by BNR fungus on decreasing proline content, but increase to height, fresh weight and dry weight. We can see without goat manure and husk charcoal dosage charcoal (P0S0). the highest proline content but height, fresh and weight of plant lowest, while in the treatment interaction (P2S3) proline content  $0.77 \text{ mol g}^{-1}$  was not significantly different from (P1S3) and (P3S3), height (113.83cm), dry weight (6.73 g) and fresh weight (1.47 g) were not significantly different from (P1S3) on vanilla, while the observed levels of proline decreased without manure treatment and husk charcoal (P0S0).

The increase in proline indicates the plant is able to survive in times of drought (Daryanti et al., 2014). Drought in vanilla has an effect on physiology and anatomy, because drought causes a decrease in the rate of photosynthesis and leaf area, and even closure of leaf stomata, at the cellular level there is an increase in the concentration of dissolved matter including plant proline (Prihastati, 2012) (Subantoro, 2014) (Siregar, 2015) (Pardosi, 2021). BNR which acts as mycorrhizae forms external hyphae that absorb nutrients around the roots

and are stored in peloton in plant tissues through internal hyphae. The peloton will be destroyed in the plant after 2 weeks, because hypha-infected tissue becomes lysed so that the plant tissue contains nutrients (Rasmussen & Rasmussen, 2009). This is supported by research conducted by (Haryuni. et al., 2020) which states that the application of biopesticides and *Fusarium* inoculation after induction of BNR can increase nitrogen, phosphorus, and potassium content of vanilla plants. Another study by (Muslim, 2019) showed that the success of biological control of fusarium wilt in tomato plants was by using hypovirulent binucleate rhizoctonia (HBNR). Another study using antagonistic fungi was carried out on *Trichoderma* sp inoculated treatment 25g/plant and water heating for 60 minutes on sugarcane bud chips seedling growth (*Saccharum officinarum*) showed the lowest proline content of 1.422 mol g<sup>-1</sup>, this caused the seeds to be drought-resistant and not easy to die (Haryuni, 2015 a).



Figure 1. Application BNR to vanilla plants at Rejosari village, Jambu district, Semarang Regency, Central of Java.

The novelty in this research is the use of husk charcoal and goat manure which is induced by BNR to control plant proline, the decrease in proline indicates that the plant is in sufficient water conditions. They really help the decomposition process that produces plant nutrients. The results of this study helped vanilla agronomists adjust the dose of husk charcoal and goat manure by influencing the BNR fungus so that plant growth was better and resistant to drought stress conditions.

## CONCLUSION

The effect of doses of goat manure and husk charcoal induced by BNR can reduce to proline and increase vanilla plant growth. The decrease in proline affects the resistance of vanilla plants to drought. The results showed that the combination of treatments (POS0), namely the dose of goat manure 0g/plant (P0) and husk choarcoal 0g/plant (S0) the plant proline was 3.94  $\mu\text{mol g}^{-1}$ , plant height 22.33 cm, fresh weight 5.20 g/plant. plant and dry

weight 1.05g/plant, while the treatment (P2S3), namely the dose of goat manure 10g/plant (P2) and husk charcoal 15 g/plant (S3), the plant proline was 0.77  $\mu\text{mol g}^{-1}$ , plant height 113.83 cm, fresh weight 6.73 g/plant. plant and dry weight 1.47g/plant.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge the research grants from Universitas Tunas Pembangunan Surakarta No. 0017/PK-P/LPPM UTP/XII-2021

## REFERENCES

- Assefa, S., & Tadesse, S. (2019). The Principal Role of Organic Fertilizer on Soil Properties and Agricultural Productivity -A Review. *Agri Res & Tech: Open Access J.*, 22(2), 556192.
- Bates, L. S., Waldren, R. P., & Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39(1), 205–207.
- Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M. I., Ashraf, M., Ahmed, N., & Zhang, L. (2019). Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance . In *Frontiers in Plant Science* (Vol. 10, p. 1068).  
<https://www.frontiersin.org/article/10.3389/fpls.2019.01068>
- Boutasknit, A., Baslam, M., Ait-El-Mokhtar, M., Anli, M., Ben-Laouane, R., Douira, A., El Modafar, C., Mitsui, T., Wahbi, S., & Meddich, A. (2020). Arbuscular Mycorrhizal Fungi Mediate Drought Tolerance and Recovery in Two Contrasting Carob (*Ceratonia siliqua* L.) Ecotypes by Regulating Stomatal, Water Relations, and (In)Organic Adjustments. *Plants*, 9(1).  
<https://doi.org/10.3390/plants9010080>
- Chun, S. C., Paramasivan, M., & Chandrasekaran, M. (2018). Proline accumulation influenced by osmotic stress in arbuscular mycorrhizal symbiotic plants. *Frontiers in Microbiology*, 9, 2525.
- Daryanti, D., Haryuni, H., & Utami, D. S. (2014). Pengimbasan Ketahanan Rhizoctonia Binukleat Terhadap Cekaman air Pada Bibit Vanili (*Vanilla Planifolia* Andrews). *Jurnal Ilmiah Agrineca*, 14(1).
- Fileccia, T., Hovhera, V. ., & Bernoux, M. (2014). *Ukraine: Soil fertility to strengthen climate resilience preliminary assessment of the potential benefits of conservation agriculture*. Food and Agriculture Organization of the United Nations.
- Frenkel, D. H., & Belanger, F. . (2018). *Handbook Of Vanilla Science And Technology*. *Food Science*

*And Technology 2nd, Wiley Blackwell.*

- Furlan, A. L., Bianucci, E., Giordano, W., Castro, S., & Becker, D. F. (2020). Proline metabolic dynamics and implications in drought tolerance of peanut plants. *Plant Physiology and Biochemistry*, *151*, 566–578. <https://doi.org/https://doi.org/10.1016/j.plaphy.2020.04.010>
- Handajaningsih, M., Marwanto, T., Armitasari, E., Inorihah., & Pujiwati, H. (2018). Growth and yield response of melon to different time application of goat manure and dose of potassium. *Akta Agrosia*, *21*(1), 1–5.
- Harahap, F. ., & Hidayat, B. (2021). The effect of rice husk ash and goat manure on the growth of spinach (*Amaranthus gangetius*). *Jurnal Pertanian Tropik*, *8*(1), 26–32.
- Haryuni., Harahap, A. F. P., Priyatmojo, A., & Gozan, M. (2020). The Effects of Biopesticide and *Fusarium oxysporum* f. sp. *vanillae* on the Nutrient Content of Binucleate *Rhizoctonia*-Induced Vanilla Plant. *International Journal of Agronomy*, 2020.
- Haryuni. (2015). The Effect of Hot Water Treatment and Dose *Trichoderma* sp. to Plant Tissue of Seedling Growth from Bud Chips of Sugarcane (*Saccharum officinarum*). *Biosaintifika Journal of Biology & Biology Education*, *7*(1), 1–7. <https://doi.org/10.15294/biosaintifika.v7i1.3541>
- Haryuni, H., Dewi, T. S. K., & Nuryati, T. (2015). Pengaruh Dosis *Rhizoctonia* Binukleat (BNR) dan Pupuk posfor terhadap Pertumbuhan benih Vanili (*Vanilla planifolia* Andrew). *Prosiding Seminar Nasional & Internasional*, 36–45.
- Haryuni, H., Harahap, A. F. P., Supartini, Priyatmojo, A., & Gozan, M. (2020). The Effects of Biopesticide and *Fusarium oxysporum* f.sp. *vanillae* on the Nutrient Content of Binucleate *Rhizoctonia* -Induced Vanilla Plant . *International Journal of Agronomy*, 2020, 1–6. <https://doi.org/10.1155/2020/5092893>
- Haryuni, H., Suprapti, E., Dewi, T. S. K., Supriyadi, T., Nugroho, A. A., Priyatmojo, A., & Gozan, M. (2018). Phosphorus dosage and cow urine to chlorophyll and proline contents on binucleate *Rhizoctonia* by induced resistance of vanilla. *Proceedings of the International Conference on Science and Education and Technology*, *247*, 215–218.
- Hendrati, R. L., Rachmawati, D., & Pamuji, A. C. (2016). Drought responses on growth, proline contents and root anatomy of *Acacia auriculiformis* Cunn., *Tectona grandis* L., *Alstonia spectabilis* Br., and *Cedrela odorata* L. *Jurnal Penelitian Kehutanan Wallacea*, *5*(2), 123–133.
- Huang, W., Ratkowsky, D. A., Hui, C., Wang, P., Su, J., & Shi, P. (2019). Leaf Fresh Weight Versus Dry Weight: Which is Better for Describing the Scaling Relationship between Leaf Biomass and Leaf Area for Broad-Leaved Plants? *Forests*, *10*(3). <https://doi.org/10.3390/f10030256>
- Hughes, A. P., Cockshull, K. E., & Heath, O. V. S. (1970). Leaf Area and Absolute Leaf Water

- Content. *Annals of Botany*, 34(2), 259–266. <https://doi.org/10.1093/oxfordjournals.aob.a084366>
- Ichsan, M. C., Santoso, I., & Oktarina, O. (2016). Uji Efektivitas Waktu Aplikasi Bahan Organik dan Dosis Pupuk Sp-36 dalam Meningkatkan Produksi Okra (*Abelmoschus Esculentus*). *Agritrop: Jurnal Ilmu-Ilmu Pertanian (Journal of Agricultural Science)*, 14(2).
- Kavi Kishor, P. B., Hima Kumari, P., Sunita, M. S. L., & Sreenivasulu, N. (2015). Role of proline in cell wall synthesis and plant development and its implications in plant ontogeny. *Frontiers in Plant Science*, 6. <https://doi.org/10.3389/fpls.2015.00544>
- Kumar, M., Kumar Patel, M., Kumar, N., Bajpai, A. B., & Siddique, K. H. M. (2021). Metabolomics and Molecular Approaches Reveal Drought Stress Tolerance in Plants. *International Journal of Molecular Sciences*, 22(17). <https://doi.org/10.3390/ijms22179108>
- Lin, S., Shao, L., Hui, C., Song, Y., Reddy, G. V. P., Gielis, J., Li, F., Ding, Y., Wei, Q., & Shi, P. (2018). Why Does Not the Leaf Weight-Area Allometry of Bamboos Follow the 3/2-Power Law? *Frontiers in Plant Science*, 9. <https://doi.org/10.3389/fpls.2018.00583>
- Muslim, A. (2019). *Pengendalian Hayati Patogen Tanaman Dengan Mikroorganisme Antagonis*. Unsri Press. <https://repository.unsri.ac.id/8930/>
- Nurcholis. (2017). Perbanyak tanaman panili (*Vanilla planifolia* Andrews) secara stek dan upaya untuk mendukung keberhasilan serta pertumbuhannya. *Agrovigor*, 10(2), 149–156.
- Osakabe, Y., Osakabe, K., Shinozaki, K., & Tran, L.-S. P. (2014). Response of plants to water stress. *Frontiers in Plant Science*, 5, 86.
- Pardosi, D. (2021). *Penampilan Morfologi Beberapa Varietas Tanaman Cabai Rawit (*Capsicum frutescens* L.) terhadap Cekaman Kekeringan*.
- Pratomo & Prasetyo, B. (2018). Pembuatan pupuk kompos berbahan feses kambing menggunakan bantuan effective microorganism (EM4), Di Desa Tegal, Bogor. *Prosiding PKM-CSR*, 1(2018), 403–412.
- Prihastati, E. (2012). Peranan dan Pola Akumulasi Proline Tanaman pada Adaptasi Cekaman Kekeringan. *Prosiding Seminar Biologi*, 9(1), 594–597.
- Rasmussen, H. N., & Rasmussen, F. N. (2009). Orchid Mycorrhiza: Implications of a Mycophagous Life Style. *Oikos*, 118(3), 334–345. <http://www.jstor.org/stable/40235687>
- Singh, R. P. (2012). Organic fertilizers: Types, production and environmental impact. In *Organic Fertilizers: Types, Production and Environmental Impact* (Issue July 2012).
- Siregar, B. L. (2015). Prolin: Metabolisme Dan Peranannya Dalam Tanaman Stress Kekeringan. *Jurnal Penelitian Ilmu Pertanian Methadagro*, 1(1), 1–8. [https://uhn.ac.id/content.php?ref=TXgL&ssid=3\\_210204112326](https://uhn.ac.id/content.php?ref=TXgL&ssid=3_210204112326)

- Subantoro, R. (2014). Pengaruh cekaman kekeringan terhadap respon fisiologis perkecambahan benih kacang tanah (*Arachis hypogaea* L). *Mediagro*, 10(2).
- Surdianto, Y., Sutrisna, N., Basuno., & Solihi. (2015). Panduan teknis cara membuat arang sekam padi. *Balai Pengkajian Teknologi Pertanian (BPTP) Jawa Barat*, 23.  
<http://repository.pertanian.go.id/handle/123456789/6751>
- Tapwal, A., Kumar, R., & Borah, D. (2015). Effect of mycorrhizal inoculations on the growth of *Shorea robusta* seedlings. *Nusantara Bioscienc*, 7(1), 1–5.
- Valadares, R. B. S., Perotto, S., Santos, E. C., & Lambais, M. R. (2014). Proteome changes in *Oncidium sphacelatum* (Orchidaceae) at different trophic stages of symbiotic germination. *Mycorrhiza*, 24(5), 349–360. <https://doi.org/10.1007/s00572-013-0547-2>
- Widiastuti, E., & Latifah, E. (2016). Keragaan Pertumbuhan dan Biomassa Varietas Kedelai (*Glycine max* (L)) di Lahan Sawah dengan Aplikasi Pupuk Organik Cair. *Jurnal Ilmu Pertanian Indonesia (JIPI)*, 21(2), 90–97.
- Yenni, M., Ibrahim, M. H., Nulit, R., & Sakimin, S. Z. (2021). The interactive effects of fertilizer and water stress on plant growth, leaf gas exchange and nutrient uptake on strawberry (*Fragaria x ananassa*, Duch). *AIMS Environmental Science*, 8(6), 597–618.
- Zhang, S., Wang, L., Ma, F., Zhang, X., & Fu, D. (2016). Arbuscular mycorrhiza improved phosphorus efficiency in paddy fields. *Ecological Engineering*, 95, 64–72.  
<https://doi.org/https://doi.org/10.1016/j.ecoleng.2016.06.029>



Home > User > Author > Submissions > #35618 > Summary

## #35618 Summary

SUMMARY REVIEW EDITING

### Submission

Authors	Haryuni Haryuni, Muhamad Amin, Endang Suprapti, Tyas Soemarah Kurnia Dewi, Eko Hartoyo
Title	Proline Reduction and Increasing Growth of Vanilla Plants Induced by BNR Fungus with Dose of Goat Manure and Husk Charcoal
Original file	35618-90509-2-SM.DOCX 2022-03-14
Supp. files	35618-90510-2-SP.DOCX 2022-03-14
Submitter	Haryuni Haryuni Haryuni
Date submitted	March 14, 2022 - 05:24 PM
Section	Articles
Editor	Dyah Indriyanti
Abstract Views	352

### Status

Status	Published Vol 14, No 1 (2022): April 2022
Initiated	2022-04-06
Last modified	2022-04-13

### Submission Metadata

#### Authors

Name	Haryuni Haryuni
ORCID iD	<a href="https://orcid.org/0000-0002-2574-5506">https://orcid.org/0000-0002-2574-5506</a>
Affiliation	Department of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta
Country	Indonesia
Bio Statement	—
Principal contact for editorial correspondence.	
Name	Muhamad Amin
Affiliation	Department of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta
Country	Indonesia
Bio Statement	—
Name	Endang Suprapti
Affiliation	Department of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta
Country	—
Bio Statement	—
Name	Tyas Soemarah Kurnia Dewi
Affiliation	Department of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta
Country	Indonesia
Bio Statement	—
Name	Eko Hartoyo
ORCID iD	<a href="https://orcid.org/0000-0003-2358-8452">https://orcid.org/0000-0003-2358-8452</a>
Affiliation	Department of Agribusiness, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta
Country	Indonesia
Bio Statement	—

#### Title and Abstract

Title	Proline Reduction and Increasing Growth of Vanilla Plants Induced by BNR Fungus with Dose of Goat Manure and Husk Charcoal
Abstract	Goat manure and husk charcoal are agricultural wastes that have not been used optimally, contain soil microorganisms that are effective in converting nutrients into more perfect forms, play a role in improving soil physical properties and increase the ability of soil to bind nutrients. The purpose of this study is to determine the effect of goat manure and husk charcoal to proline reduction and increase growth of vanilla plant induced by Binucleate Rhizoctonia (BNR). The study was designed with a



#### ABOUT THE JOURNAL

Aims and Scope

Publication Ethics

Indexing & Abstracting

Editorial Team

Contact

#### COLLABORATE WITH



#### FOR AUTHORS

Guidelines for Author

Manuscript Template

Mendeley Guidelines

Scopus Citation Analysis

#### Readers

ID 328,827	PK 1,146
US 21,887	VN 1,059
CN 11,072	IR 1,008
PH 10,460	NG 1,007
IN 6,316	GB 975
MY 4,142	MX 825
SG 2,693	AU 802
TH 2,256	KR 796
BR 1,606	CA 740
JP 1,406	EG 737
TR 1,169	TW 656

Pageviews: 1,077,522

Flags Collected: 198



View My Stats

#### USER

You are logged in as...

**haryuni**

- » My Journals
- » My Profile
- » Log Out



BNR can reduce proline contents and increase growth of vanilla plant. The benefit of this research is to provide treatment information to reduce proline levels and increase plant growth with the application of goat fertilizer and husk charcoal.

## Indexing

Keywords binucleate-Rizoctonia, husk, manure, vanilla  
Language en

## Supporting Agencies

Agencies Universitas Tunas Pembangunan

## OpenAIRE Specific Metadata

ProjectID —

## References

- Assefa, S., & Tadesse, S. (2019). The Principal Role of Organic Fertilizer on Soil Properties and Agricultural Productivity -A Review. *Agri Res & Tech: Open Access J.*, 22(2), 556192.
- Bates, L. S., Waldren, R. P., & Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39(1), 205–207.
- Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M. I., Ashraf, M., Ahmed, N., & Zhang, L. (2019). Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance. In *Frontiers in Plant Science* (Vol. 10, p. 1068). <https://www.frontiersin.org/article/10.3389/fpls.2019.01068>
- Boutasknit, A., Baslam, M., Ait-El-Mokhtar, M., Anli, M., Ben-Laouane, R., Douira, A., El Modafar, C., Mitsui, T., Wahbi, S., & Meddich, A. (2020). Arbuscular Mycorrhizal Fungi Mediate Drought Tolerance and Recovery in Two Contrasting Carob (*Ceratonia siliqua* L.) Ecotypes by Regulating Stomatal, Water Relations, and (In)Organic Adjustments. *Plants*, 9(1). <https://doi.org/10.3390/plants9010080>
- Chun, S. C., Paramasivan, M., & Chandrasekaran, M. (2018). Proline accumulation influenced by osmotic stress in arbuscular mycorrhizal symbiotic plants. *Frontiers in Microbiology*, 9, 2525.
- Daryanti, D., Haryuni, H., & Utami, D. S. (2014). Pengimbasan Ketahanan Rhizoctonia Binukleat Terhadap Cekaman air Pada Bibit Vanili (*Vanilla planifolia* Andrews). *Jurnal Ilmiah Agrinca*, 14(1).
- Fileccia, T., Hovhera, V., & Bernoux, M. (2014). Ukraine: Soil fertility to strengthen climate resilience preliminary assessment of the potential benefits of conservation agriculture. Food and Agriculture Organization of the United Nations.
- Frenkel, D. H., & Belanger, F. (2018). *Handbook Of Vanilla Science And Technology*. Food Science And Technology 2nd, Wiley Blackwell.
- Furlan, A. L., Bianucci, E., Giordano, W., Castro, S., & Becker, D. F. (2020). Proline metabolic dynamics and implications in drought tolerance of peanut plants. *Plant Physiology and Biochemistry*, 151, 566–578. <https://doi.org/10.1016/j.plaphy.2020.04.010>
- Handajaningsih, M., Marwanto, T., Armitasari, E., Inorihah, & Pujiwati, H. (2018). Growth and yield response of melon to different time application of goat manure and dose of potassium. *Akta Agrosia*, 21(1), 1–5.
- Harahap, F., & Hidayat, B. (2021). The effect of rice husk ash and goat manure on the growth of spinach (*Amaranthus gangeticus*). *Jurnal Pertanian Tropik*, 8(1), 26–32.
- Haryuni, Harahap, A. F. P., Priyatmojo, A., & Gozan, M. (2020). The Effects of Biopesticide and *Fusarium oxysporum* f. sp. *vanillae* on the Nutrient Content of Binucleate *Rhizoctonia*-Induced Vanilla Plant. *International Journal of Agronomy*, 2020.
- Haryuni. (2015). The Effect of Hot Water Treatment and Dose *Trichoderma* sp. to Plant Tissue of Seedling Growth from Bud Chips of Sugarcane (*Saccharum officinarum*). *Biosaintifika Journal of Biology & Biology Education*, 7(1), 1–7. <https://doi.org/10.15294/biosaintifika.v7i1.3541>
- Haryuni, H., Dewi, T. S. K., & Nuryati, T. (2015). Pengaruh Dosis *Rhizoctonia* Binukleat (BNR) dan Pupuk posfor terhadap Pertumbuhan benih Vanili (*Vanilla planifolia* Andrew). *Prosiding Seminar Nasional & Internasional*, 36–45.
- Haryuni, H., Harahap, A. F. P., Supartini, Priyatmojo, A., & Gozan, M. (2020). The Effects of Biopesticide and *Fusarium oxysporum* f. sp. *vanillae* on the Nutrient Content of Binucleate *Rhizoctonia* -Induced Vanilla Plant. *International Journal of Agronomy*, 2020, 1–6. <https://doi.org/10.1155/2020/5092893>
- Haryuni, H., Suprapti, E., Dewi, T. S. K., Supriyadi, T., Nugroho, A. A., Priyatmojo, A., & Gozan, M. (2018). Phosphorus dosage and cow urine to chlorophyll and proline contents on binucleate *Rhizoctonia* by induced resistance of vanilla. *Proceedings of the International Conference on Science and Education and Technology*, 247, 215–218.
- Hendrati, R. L., Rachmawati, D., & Pamuji, A. C. (2016). Drought responses on growth, proline contents and root anatomy of *Acacia auriculiformis* Cunn., *Tectona grandis* L., *Alstonia spectabilis* Br., and *Cedrela odorata* L. *Jurnal Penelitian Kehutanan Wallacea*, 5(2), 123–133.
- Huang, W., Ratkowsky, D. A., Hui, C., Wang, P., Su, J., & Shi, P. (2019). Leaf Fresh Weight Versus Dry Weight: Which is Better for Describing the Scaling Relationship between Leaf Biomass and Leaf Area for Broad-Leaved Plants? *Forests*, 10(3). <https://doi.org/10.3390/f10030256>
- Hughes, A. P., Cockshull, K. E., & Heath, O. V. S. (1970). Leaf Area and Absolute Leaf Water Content. *Annals of Botany*, 34(2), 259–266. <https://doi.org/10.1093/oxfordjournals.aob.a084366>
- Ichsan, M. C., Santoso, I., & Oktarina, O. (2016). Uji Efektivitas Waktu Aplikasi Bahan Organik dan Dosis Pupuk Sp-36 dalam Meningkatkan Produksi Okra (*Abelmoschus Esculentus*). *Agritrop: Jurnal Ilmu-Ilmu Pertanian (Journal of Agricultural Science)*, 14(2).
- Kavi Kishor, P. B., Hima Kumari, P., Sunita, M. S. L., & Sreenivasulu, N. (2015). Role of proline in cell wall synthesis and plant development and its implications in plant ontogeny. *Frontiers in Plant Science*, 6. <https://doi.org/10.3389/fpls.2015.00544>
- Kumar, M., Kumar Patel, M., Kumar, N., Bajpai, A. B., & Siddique, K. H. M. (2021). Metabolomics and Molecular Approaches Reveal Drought Stress Tolerance in Plants. *International Journal of Molecular Sciences*, 22(17). <https://doi.org/10.3390/ijms22179108>
- Lin, S., Shao, L., Hui, C., Song, Y., Reddy, G. V. P., Gielis, J., Li, F., Ding, Y., Wei, Q., & Shi, P. (2018). Why Does Not the Leaf Weight-Area Allometry of Bamboos Follow the 3/2-Power Law? *Frontiers in Plant Science*, 9. <https://doi.org/10.3389/fpls.2018.00583>
- Muslim, A. (2019). *Pengendalian Hayati Patogen Tanaman Dengan Mikroorganisme Antagonis*. Unsri Press. <https://repository.unsri.ac.id/8930/>
- Nurcholis. (2017). Perbanyak tanaman panili (*Vanilla planifolia* Andrews) secara stek dan upaya untuk mendukung keberhasilan serta pertumbuhannya. *Agrovigor*, 10(2), 149–156.
- Osakabe, Y., Osakabe, K., Shinozaki, K., & Tran, L.-S. P. (2014). Response of plants to water stress. *Frontiers in Plant Science*, 5, 86.
- Pardosi, D. (2021). Penampilan Morfologi Beberapa Varietas Tanaman Cabai Rawit (*Capsicum frutescens* L.) terhadap Cekaman Kekeringan.
- Pratomo & Prasetyo, B. (2018). Pembuatan pupuk kompos berbahan feses kambing menggunakan bantuan effective

- Rasmussen, H. N., & Rasmussen, F. N. (2009). Orchid Mycorrhiza: Implications of a Mycophagous Life Style. *Oikos*, 118(3), 334–345. <http://www.jstor.org/stable/40235687>
- Singh, R. P. (2012). Organic fertilizers: Types, production and environmental impact. In *Organic Fertilizers: Types, Production and Environmental Impact* (Issue July 2012).
- Siregar, B. L. (2015). Prolin: Metabolisme Dan Perannya Dalam Tanaman Stress Kekeringan. *Jurnal Penelitian Ilmu Pertanian Methadagro*, 1(1), 1–8. [https://uhn.ac.id/content.php?ref=T XgL&ssid=3\\_210204112326](https://uhn.ac.id/content.php?ref=T XgL&ssid=3_210204112326)
- Subantoro, R. (2014). Pengaruh cekaman kekeringan terhadap respon fisiologis perkecambahan benih kacang tanah (*Arachis hypogaea* L.). *Medi agro*, 10(2).
- Surdianto, Y., Sutrisna, N., Basuno, & Solihi. (2015). Panduan teknis cara membuat arang sekam pa di. Balai Pengkajian Teknologi Pertanian (BPTP) Jawa Barat, 23. <http://repository.pertanian.go.id/handle/123456789/6751>
- Tapwal, A., Kumar, R., & Borah, D. (2015). Effect of mycorrhizal inoculations on the growth of *Shorea robusta* seedlings. *Nusantara Bioscien*, 7(1), 1–5.
- Valadares, R. B. S., Perotto, S., Santos, E. C., & Lambais, M. R. (2014). Proteome changes in *Oncidium sphacelatum* (Orchidaceae) at different trophic stages of symbiotic germination. *Mycorrhiza*, 24(5), 349–360. <https://doi.org/10.1007/s00572-013-0547-2>
- Widiastuti, E., & Latifah, E. (2016). Keragaan Pertumbuhan dan Biomassa Varietas Kedelai (*Glycine max* (L)) di Lahan Sawah dengan Aplikasi Pupuk Organik Cair. *Jurnal Ilmu Pertanian Indonesia (JIPI)*, 21(2), 90–97.
- Yenni, M., Ibrahim, M. H., Nulit, R., & Sakimin, S. Z. (2021). The interactive effects of fertilizer and water stress on plant growth, leaf gas exchange and nutrient uptake on strawberry (*Fragaria x ananassa*, Duch). *AIMS Environmental Science*, 8(6), 597–618.
- Zhang, S., Wang, L., Ma, F., Zhang, X., & Fu, D. (2016). Arbuscular mycorrhiza improved phosphorus efficiency in paddy fields. *Ecological Engineering*, 95, 64–72. <https://doi.org/10.1016/j.ecoleng.2016.06.029>



This work is licensed under a Creative Commons Attribution 4.0 International License.



Home > User > Author > Submissions > #35618 > Review

## #35618 Review

SUMMARY REVIEW EDITING

### Submission

Authors Haryuni Haryuni, Muhamad Amin, Endang Suprapti, Tyas Soemarah Kurnia Dewi, Eko Hartoyo

Title Proline Reduction and Increasing Growth of Vanilla Plants Induced by BNR Fungus with Dose of Goat Manure and Husk Charcoal

Section Articles

Editor Dyah Indriyanti

### Peer Review

#### Round 1

Review Version 35618-90511-1-RV.DOCX 2022-03-14

Initiated 2022-03-28

Last modified 2022-03-28

Uploaded file None

### Editor Decision

Decision Accept Submission 2022-04-05

Notify Editor Editor/Author Email Record No Comments

Editor Version 35618-91136-1-ED.DOCX 2022-03-28

Author Version None

Upload Author Version  No file chosen



This work is licensed under a Creative Commons Attribution 4.0 International License.



#### ABOUT THE JOURNAL

- Aims and Scope
- Publication Ethics
- Indexing & Abstracting
- Editorial Team
- Contact

#### COLLABORATE WITH



#### FOR AUTHORS

- Guidelines for Author
- Manuscript Template
- Mendeley Guidelines
- Scopus Citation Analysis

#### Readers

ID 328,827	PK 1,146
US 21,887	VN 1,059
CN 11,072	IR 1,008
PH 10,460	NG 1,007
IN 6,316	GB 975
MY 4,142	MX 825
SG 2,693	AU 802
TH 2,256	KR 796
BR 1,606	CA 740
JP 1,406	EG 737
TR 1,169	TW 656

Pageviews: 1,077,522  
Flags Collected: 198



View My Stats

#### USER

You are logged in as...

**haryuni**

- » My Journals
- » My Profile
- » Log Out

## EFFECT OF GOAT MANURE AND HUSK CHARCOAL DOSAGE TO PROLINE AND GROWTH BY BNR-INDUCED VANILLA

Haryuni<sup>1\*</sup>, Amin, M<sup>2</sup>, Rumaningsih, M<sup>3</sup>, Suprapti, E<sup>1</sup>, Tyas S. K Dewi<sup>1</sup>, Hartoyo, E<sup>4</sup>, Supartini<sup>3</sup>, Maryani, I.E<sup>3</sup>,

<sup>1</sup> Department of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta, Central of Java, Indonesia 57139

<sup>2</sup> Department student of Agronomy, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta, Central of Java, Indonesia 57139

<sup>3</sup> Department of Management, Faculty of Economy and Bisnis Universitas Tunas Pembangunan, Surakarta, Central of Java, Indonesia 57135

<sup>4</sup> Department of Agribisnis, Faculty of Agriculture, Universitas Tunas Pembangunan, Surakarta, Central of Java, Indonesia 57139

\*Corresponding author: haryuni@lecture.utp.ac.id

### Abstract

Goat manure is one of the organic fertilizers that plays a role in increasing soil pH and soil nutrient availability, while husk charcoal as agricultural waste is very large and has not been used optimally so it needs to be changed for use and in the composting process. These interactions become complex materials containing soil microorganisms that are effective in converting nutrients from soils for plants into more perfect forms. The goat manure and husk charcoal are essential for soil nutrients, improving soil physicality and increasing the ability of soil to bind nutrients.

The purpose of this study is to determine the effect of goat manure and husk charcoal to proline and grow by Binucleate Rhizoctonia (BNR) induced vanilla plant.

The study was designed with a factorial completely randomized block design. The first factor is the dose of manure (0,5,10,15) g/plant, the second factor is the dose of husk charcoal (0,5, 10, 15) g/plant.

The application of goat manure and rice husk dosage induced by BNR can decrease proline content and increase the growth of vanilla plant.

**Keywords:** binucleate-Rizoctonia, husk, manure, vanilla

### INTRODUCTION

Vanilla is one of the plantation crops which is the country's foreign exchange reserves from exports. Indonesian vanilla comes from Mexico, the first use of vanilla was done by the Aztecs as a preserved chocolate fragrance that enhances the taste. Vanillin (C<sub>8</sub>H<sub>8</sub>O<sub>3</sub>) content which causes vanilla (Nurcholis, 2017) to give off a distinctive aroma used in the

**Commented [A1]:** Judul kurang menarik mhn dibuat judul yg jelas dan menarik. Bisa dimulai dr hasil yg dicapai melalui proses...apa...  
Jika dilihat tujuan hanya ingin melihat pertumbuhan saja...mhn dicek kembali

apakah ada departemen student?, samakan saja dengan no 1, shg nomornya hanya 3 saja

Nama author yg tinggal ditulis dobel... Haryuni Haryuni...  
Kalau tdk dobel nanti jika disitasi Mendeley akan hilang namanya tinggal tanda titik.

**Commented [A2]:** Nama ditulis lengkap jangan disingkat.  
Nama satu kata ditulis ulang (dobel)

**Commented [A3]:** Latar belakang terlalu panjang cukup 2-3 baris saja mengantarkan tujuan penelitian

**Commented [A4]:** Proline ...asam amino? Maksudnya apakah kandungan asam amino?

Tujuan di abstrak tdk sama dengan tujuan diakhir pendahuluan...mhn dicek lagi, mana yg benar

**Commented [A5]:** Metode kurang lengkap ,mhn dilengkapi sampai analisis datanya apa

**Commented [A6]:** Simpulannya mhn diperjelas sesuai tujuan.  
Beri keterangan hasil penelitian ini apa?  
Apa manfaatnya?

Total abstrak maksimal 250 kata

**Commented [A7]:** Gunakan bhs inggris yg benar... to give off .. Apa maksudnya?

food, beverage, cosmetic, and pharmaceutical industries. Vanilla plants are characterized by growth that propagates up to 30 meters, epiphytes and vanilla fruit producing pods (Havkin-Frenkel, D. & Belanger, 2018). The initial growth of vanilla requires a medium that has the ability to hold water, loose soil structure, smooth aeration and drainage, does not experience drought and contains sufficient nutrients needed for growth (Fileccia et al., 2014)

Fertilization treatments, especially organic fertilizers, accelerate soil decomposition so that soil microorganisms can help provide nutrients needed by plants. Organic fertilizer derived from goat manure has the advantage of increasing soil humus content, stimulating soil microorganisms, and improving soil structure so as to increase the availability of nutrients for plants (Singh, 2012); (Assefa & Tadesse, 2019); (Pratomo, H., & Prasetyo, 2018). Application of organic matter derived from goat manure enriches microorganisms, produces nutrients (macro and micro), increases soil aggregates so as to increase water infiltration and acts as a soil enhancer. Goat manure has a long-term effect on the potassium content of 2.88%, cow manure 0.69%, chicken manure 1.45%, other nutrients nitrogen 2.77%, phosphorus 1.78% (Handajansih et al., 2018).

In addition to goat manure, soil organic matter can also be added by giving rice husk charcoal. Treatment with husk charcoal increased growth and reduced the percentage of seedling mortality caused by the fungus *Fusarium oxysporum* f. sp. *Vanilla* (Haryuni et al., 2018). The results of the decomposition of organic matter from goat manure and husk charcoal (N, P, K, Ca, S and Mg) which were eliminated directly were able to increase the pH and increase the ability of the soil as a cation buffer because the decomposition of organic matter produced a complex compound (humus) (Fileccia et al., 2014). Organic fertilizer derived from goat manure has the advantage of increasing soil humus content, stimulating soil microorganisms, and improving soil structure so as to increase the availability of nutrients for plants. Continuous use of land without being followed by nutrient supply, it is difficult to obtain maximum production considering the very limited capacity of the soil, the provision of goat manure with an appropriate dose is expected to help the availability of soil nutrients, goat manure and rice husk ash serve as biofertilizers (Harahap & Hidayat, 2021).

Plant growth is influenced by the availability of nutrients that can be utilized by roots and available to plants, this role can be carried out by Binucleate Rhizoctonia (BNR). The pelotoon that is in the root tissue plays a role in storing nutrients that are absorbed by external hyphae and sent into plant cells (Haryuni, et al., 2020). The material in the peloton that is in the network is utilized by plants during the metabolic process. In conditions of water

**Commented [A8]:** Pelajari cara penulisan referensi yg benar.. Gunakan satu kata nama terakhir,

**Commented [A9]:** ;

**Commented [A10]:** .

**Commented [A11]:** Cek bhs inggris yg benar... Gunakan grammer yg standar

**Commented [A12]:** Apa maksudnya rice husk? Gunakan kata yg benar

**Commented [A13]:** Apa arti peloton?

shortage, the material contributes to the availability of water for plants (Begum et al., 2019).  
The interconnected cellular processes of BNR and BNR-mediated mechanisms regulate plant tolerance to drought and promote plant improvement in bio-amelioration of stresses (Boutasknit et al., 2020)

Proline is an indicator of a plant's ability to maintain turgor through a decrease in osmotic potential so that an increase in proline content indicates an increase in plant tolerance to water shortages, which results in plant growth and the quantity and quality of crop yields. Proline affects the air potential, opening and closing of leaf stomata further down regulates photosynthesis and CO<sub>2</sub> deficiency causes plant stress (Chun et al., 2018); (Osakabe et al., 2014). There is not enough information on the application of goat manure and husk charcoal to BNR-induced vanilla plants and their effect on plant proline levels.

This study aimed to determine the effect and interaction of goat manure and husk charcoal on the growth of vanilla induced by the BNR.

Jelaskan keterbaruan hasil penelitian ini, jelaskan apa manfaat penelitian ini

## MATERIAL AND METHODS

The research was carried out at Rejosari village, Jambu district, Semarang Regency, Central of Java from March to September 2018, the altitude is around 800-900 m above sea level, soil type andosol soil, each treatment was repeated 5 times used samples. Proline analysis on Laboratory of fisiology at Universitas Gadjah Mada Yogyakarta.

### Research design

Vanilla plants that are 2 years old in the field, sprinkled with BNR 10 g/plant around the plant roots by covering the surface soil, after 2 weeks of giving BNR goat manure is given at a dose of 0.5,10,15 g/plant as (P0 , P1, P2, P3) every 2 weeks up to 4 times. The second factor was the dose of husk charcoal 0,5,10,15 g/plant as (S0, S1, S2, S3) added every 2 weeks up to 4 times.

### Preparation of goat manure

The manure used is derived from wind-dried goat manure until the moisture content reaches 60%, put in a wooden box then added with molasses/Javanese sugar and coconut water in a ratio (10:1:3) mixed evenly and stored closed, every 7 days once inverted and

Commented [A14]: Kalimat ini apa maksudnya?

Commented [A15]: Proline itu senyawa apa? Apakah asam amino? Jelaskan, jelaskan keterkaitan proline dengan BNR

Commented [A16]: Pada tujuan ini tdk ada kata prolannya? Dari indtroduction mhn disusun kembali paragraf secara sistematis sesuai tujuan penelitian, kaitkan antara tujuan satu dengan yg lain, beri referensi terkait dari jurnal penelitian.

Commented [A17]: Jika dicantumkan th nya, maka penelitian ini sdh lama 4 th lalu, sebaiknya tdk ditulis th nya hanya ditulis musim atau bulan nyan saja

Commented [A18]: Jelaskan metodenya secara rinci, tempatnya bukan disini, buat metode secara sistematis dari awal hingga analisis data yg jelas.

Commented [A19]: Pindah jangan disini

Commented [A20]: Mhn dibuat metode yg sistematis tahap demi tahap spy jelas..cek semua paragraf di metode

Commented [A21]: Harusnya ini dulu ya? Jelaskan step by step metode yg dilakukan. Perlakuan goat manure dan arang sekam apakah dicampur atau sendiri sendiri?

stirred for 5 times. If the fertilizer has changed color and comes off when held by hand, the fertilizer is ready to use (Pratomo, H., & Prasetyo, 2018).

#### **Preparation of rice husk charcoal**

The rice bran is cleaned of other impurities and collected together and then a chimney/tube is installed in the middle which is made of perforated wire. wire is shaped like a tube as a means of combustion. This means that the flames will stick out. For burning, it is better to use a base made of zinc plate placed on a flat, hard and heat-resistant floor. The hole in the chimney is filled with charcoal and doused with kerosene, added with paper, so that a fire arises, when it is black the unburned part is piled up and moved near the chimney. After everything is burned, the husks turn black, allow to cool and are ready to be used for planting media mixtures (Surdianto et al., 2015)

#### **Preparation of Binucleat Rhizoctonia (BNR)**

The BNR used was as in the procedure (Haryuni et al., 2015 b) but only lasted about 1-2 months, so it was propagated in kaolin flour medium. This gives the BNR a longer shelf life of about 6 months when stored at room temperature.

#### **Observation of Plant Growth Vanilla**

Observations of plant growth were carried out 4 weeks after the first treatment of goat manure and husk charcoal including, plant height (measured from the base of the stem to the top of the stem), number of leaves (counted for all the leaves that grew for each sample plant), leaf fresh weight (taken 10 leaves). leaf starting with 5 leaves from the stem surface and then weighed), leaf dry weight all leaves were observed on leaf fresh weight by drying the leaves and continued in an oven at 105°C for 2 h and continuously at 80°C until the weight was constant (Huang et al., 2019).

#### **Measurement of Proline Content**

Analysis of proline content in vanilla plants was determined following (Bates et al., 1973).

#### **Statistical analysis**

The data obtained from the observations were analyzed by means of variance (Anova). If significantly different, Duncan's Multiple Range Test (DMRT) test at 5% level was carried out, to distinguish between treatments. To find out the direct and indirect relationship, regression and correlation analysis were carried out.

**Commented [A22]:** Apakah benar bhs inggris ini? Atau bhs indonesia yg diinggriskan? Kok aneh ada rice = nasi ?? Mhn bhs dikonsultasikan pd ahlinya

**Commented [A23]:** Setelah menyiapkan lalu bagaimana perlakuannya pd tanaman? Jelaskan langkah langkahnya.. Baru setelah itu dilakukan pengamatan (paragraf dibawah ini)

**Commented [A24]:** Jelaskan bagiana analisis prolin, bagian apanya yg dianalisis dimana... hasilnya bagaimana dll lalu bagaimana dengan pengamatan pertumbuhan atau growth ?

**Commented [A25]:** Data apa yg dianalisis? Jelaskan,

## RESULT AND DISCUSSION

Commented [A26]: Beri foto hasil penelitian terbaik

**Sebelum mencantumkan tabel beri kalimat pengantar dulu jangan tiba tiba tabel...**

Table 1. Effect of goat manure on proline content, height, fresh weight and dry weight of vanilla plants induced by BNR fungus.

Treatment	Observation of parameters			
	Proline content ( $\mu\text{mol g}^{-1}$ )	height of plant (cm)	fresh weight of plant (g)	dry weight of plant (g)
P0	2.12 b	50.17 a	6.05 b	1.27 c
P1	2.06 b	58.71 ab	6.14 b	1.31 b
P2	1.91 ab	64.88 bc	6.19 ab	1.34 a
P3	1.77 a	71.08 c	6.37 a	1.36 a

Commented [A27]: Penulisan desimal salah, cek semua penulisan desimal dlm bhs inggris

Note: Numbers followed by the same letter show no significant difference in the DMRT test at 5% level.

Table 1 shows that goat manure has a significant effect on proline content, height, fresh weight, and dry weight of vanilla plant (Tabel 1). It caused by goat manure contains macro and micro nutrients rich in phosphorus, nitrogen and potassium which are needed for plant growth and development. Increasing the dose of goat manure decreased and significantly different on plant proline content (Yenni et al., 2021), are (P0)  $2.21 \mu\text{mol g}^{-1}$ , (P3)  $1.77 \mu\text{mol g}^{-1}$ . But it tends to increase and significantly different on height of vanilla plant (P0) 50.17cm, (P3) 71.08cm. the fresh weight of plant (P0) was 6.05 g, (P3) 6.37g, and the dry weight of plant (P0) was 1.27 g, (P3) 1.36 g. There are caused by BNR fungi on plants acts as mycorrhizae capable of forming internal and external hyphae on roots and peloton in plant tissues (Haryuni et al., 2020). Table 1 shows that increasing the dose of goat manure fertilizer decreased plant proline content and increased plant height, plant fresh weight, plant dry weight.

Proline is an indicator of plant stress against drought tolerance of peanut plants (Furlan et al., 2020) with the addition of goat manure to help soil microorganisms more readily utilized by plants and then stored in plant roots and stems in pelotons produced from BNR fungus inoculation. This is the same as research carried out by (Zhang et al., 2016) that BNR application as mycorrhizal can increase total root long on the black grasshopper plan. BNR fungus forms external and internal hyphae on roots and forms peloton in plant transport bundles, microorganisms contained in goat manure are able to improve soil structure so that

Commented [A28]: Jelaskan prolin itu apa? Apakah salah satu nama asam amino? Apa fungsi nya bagi tanaman?

Commented [A29]: Apa arti pelotons ?



water in the soil is more easily absorbed by plant roots. Lack of water of vanilla plant tissues is indicated by high levels of proline (Haryuni et al., 2020).

Accumulation of free **proline** as a form of general response of a plant to stress factors in drought, salinity, cold air and pathogen . Height, fresh weight and dry weight are indicators of plant growth and development of the plant, groundwater that enters through the roots is stored in plant tissue, used in metabolic processes and used for plant growth and development. Lack of water in plants will affect physiological processes and even cause plants to die (Pardosi, 2021)

Table 2. Effect of husk charcoal dose on proline content, height, fresh weight and dry weight on vanilla plants induced by BNR fungus

Treatment	Observation of parameters			
	Proline content ( $\mu\text{mol g}^{-1}$ )	height of plant (cm)	fresh weight of plant (g)	dry weight of plant (g)
S0	3.59 d	25.04 a	5.41 d	1.13 d
S1	2.14 c	37.83 b	6.13 c	1.28 c
S2	1.27 b	73.21 c	6.50 b	1.40 b
S3	0.85 a	108.75 d	6.70 a	1.47 a

Note: Numbers followed by the same letter show no significant difference in the DMRT test at 5% level.

Table 2 shows the significant effect between the application treatment of husk charcoal and without husk charcoal on proline content, plant height, weight, and dry weight of plant. Proline content were significantly different and decreased with the application of husk charcoal at a dose of 15 g/plant (S3) by  $0.85 \mu\text{mol g}^{-1}$  all (S0)  $3.59 \mu\text{mol g}^{-1}$  but tended to increase and significantly different in plant height from 25.04 cm (S0) to 108.75 (S3). the weight of the wet stove from 5.41 g (S0) to 6.70 g (S3) and the dry weight of 1.13 g (S0) to 1.47 g (S0).

The treatment of increasing doses of rice husk charcoal, proline has increased, in accordance with the opinion of (Hendrati et al., 2016), proline will increase in tolerant plants not in sensitive plants, so that proline can be used as an indicator of drought tolerant from *Acacia auriculiformis* Cunn., *Tectona grandis* L., *Alstonia spectabilis* Br., and *Cedrela odorata* L. that are susceptible to drought. resistant to drought does not produce proline. Supported by the opinion of (Ichsan et al., 2016), The role of husk charcoal as soil organic matter stimulates soil granulation, cohesion and plasticity, improves soil structure and increases soil resistance to water and increases activity and number of soil microorganisms.

The increase in soil organisms was supported by the induction of the BNR which was inoculated at the time of *Shorea robusta* seedlings (Tapwal et al., 2015).

The increase in the fresh weight of plant was indicated by an increase in the dose of rice husk charcoal, the fresh weight was related to the photosynthate of the plant that was stored in the tissues and the absorption of water through the roots, the availability of nutrients and water in the soil affected the growth and fresh weight. Dry weight is an indicator of photosynthetic accumulation of plants stored in soybean plant tissue used for growth and development (Widiastuti & Latifah, 2016). However, many **investigators** used leaf dry weight as the representative of leaf biomass when studying the scaling relationship between leaf biomass and area. The scaling of leaf mass and leaf area could directly influence the amount of light-capturing surface area, which gives an investment of dry biomass (Huang et al., 2019).

**To determine** leaf biomass related to leaf moisture content, by measuring leaf dry weight and leaf fresh weight so that it can determine water loss, to make it easier to determine gas exchange (CO<sub>2</sub>, O<sub>2</sub>, and H<sub>2</sub>O) leaf dry weight must be constant. (Hughes et al., 1970) reported that leaf area of several dicot genotypes had a linear relationship with absolute leaf water content (Lin et al., 2018) described the comparison between fresh leaf weight and leaf area of 11 bamboo species.

Table 3. The interaction effect of goat manure and husk charcoal dose on proline content, height, fresh weight and dry weight on vanilla plants induced by BNR fungus

Treatment	Observation of parameters			
	Proline content (μmol g <sup>-1</sup> )	height of plant (cm)	fresh weight of plant (g)	dry weight of plant (g)
P0S0	3.94 j	22.33 a	5.20 f	1.05 h
P1S0	3.45 hi	25.00 a	5.23 f	1.10 h
P2S0	3.65 i	25.00 a	5.33 f	1.16 g
P3S0	3.33 h	27.83 a	5.90 e	1.20 fg
P0S1	2.35 g	30.00 ab	6.00 ed	1.23 ef
P1S1	2.34 g	33.67 ab	6.13 cde	1.26 de
P2S1	2.37 g	40.33 bc	6.17 cde	1.30 cd
P3S1	1.51 f	47.33 cd	6.23 bcde	1.32 cd
P0S2	1.09 bc	54.00 d	6.40 abcd	1.34 c
P1S2	1.21 cde	73.50 e	6.50 abc	1.40 b
P2S2	1.42 ef	80.33 e	6.53 abc	1.42 b
P3S2	1.35 def	85.00 ef	6.60 ab	1.43 ab
P0S3	1.11 bcd	94.33 fg	6.63 ab	1.47 a
P1S3	0.64 a	102.67 g	6.70 a	1.47 a

**Commented [A30]:** Ini siapa? Apakah yg dimaksud peneliti? Researcher ?

**Commented [A31]:** Awali dengan subjek jangan To

**Commented [A32]:** Referensi judul mhn diganti yg terbaru

P2S3	0.77 a	113.83 h	6.73 a	1.47 a
P3S3	0.89 ab	124.17 h	6.77 a	1.47 a

Note: Numbers followed by the same letter show no significant difference in the DMRT test at 5% level.

Table 3 shows the interaction effect of goat manure and rice husk dosage charcoal induced by BNR fungus on decreasing proline content, but increase to height, fresh weight and dry weight. We can see without goat manure and rice husk dosage charcoal (POS0). the highest proline content but height, fresh and weight of plant lowest, while in the treatment interaction (P2S3) proline content  $0.77 \text{ mol g}^{-1}$  was not significantly different from (P1S3) and (P3S3), height (113.83cm), dry weight (6.73 g) and fresh weight (1.47 g) were not significantly different from (P1S3) on vanilla, while the observed levels of proline decreased without manure treatment and husk charcoal (POS0).

The increase in proline indicates the plant is able to survive in times of drought (Daryanti et al., 2014). Drought in vanilla has an effect on physiology and anatomy, because drought causes a decrease in the rate of photosynthesis and leaf area, and even closure of leaf stomata, at the cellular level there is an increase in the concentration of dissolved matter including plant proline (Prihastati, 2012) (Subantoro, 2014) (Siregar, 2015) (Pardosi, 2021). BNR which acts as mycorrhizae forms external hyphae that absorb nutrients around the roots and are stored in peloton in plant tissues through internal hyphae. The peloton will be destroyed in the plant after 2 weeks, because hypha-infected tissue becomes lysed so that the plant tissue contains nutrients (Rasmussen, 1995). This is supported by research conducted by (Haryuni. et al., 2020) which states that the application of biopesticides and Fusarium inoculation after induction of BNR can increase nitrogen, phosphorus, and potassium content of vanilla plants. Another study by (Muslim, 2019) showed that the success of biological control of fusarium wilt in tomato plants was by using hypovirulent binucleate rhizoctonia (HBNR). Another study using antagonistic fungi was carried out on *Trichoderma* sp inoculated treatment 25g/plant and water heating for 60 minutes on sugarcane bud chips seedling growth (*Saccharum officinarum*) showed the lowest proline content of  $1.422 \text{ mol g}^{-1}$ , this caused the seeds to be drought-resistant and not easy to die (Haryuni, 2015 a).

Commented [A33]: Tanda koma menandakan desimal atau ribuan? Mhn dicek

Jelaskan keterbaharuan dari penelitian ini...

Jelaskan manfaat penelitian ini

## CONCLUSION

The application of doses of goat manure and husk charcoal induced by BNR can reduce proline and increase vanilla growth. The results showed that the combination of treatments (P2S3), namely the dose of goat manure 10g/plant (P2) and husk charcoal 15 g/plant (S3), the plant proline content was 0.77  $\mu\text{mol g}^{-1}$ , plant height 113.83 cm, fresh weight 6.73 g/plant. plant and dry weight 1.47g/plant.

**Commented [A34]:** Mohon dicek dengan tujuan penelitian apakah sdh menjawab tujuan ? Cek juga dengan judul penelitian

## ACKNOWLEDGMENTS

The authors gratefully acknowledge the research grants from the Directorate General of Higher Education, Ministry of Research, Technology, and Higher Education, the Republic of Indonesia (NR: 009/K6/KM/SP2H/PENELITIAN/2018)

**Commented [A35]:** Mengingat penelitian sdh 4 th lalu sebaiknya tdk perlu di ucapan terima kasih (delete saja) krn sdh kedaluarsa 4 th lalu

## REFERENCES

- Assefa, S., & Tadesse, S. (2019). The Principal Role of Organic Fertilizer on Soil Properties and Agricultural Productivity -A Review. *Agri Res & Tech: Open Access J.*, 22(2), 556192.
- Bates, L. S., Waldren, R. P., & Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39(1), 205–207.
- Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M. I., Ashraf, M., Ahmed, N., & Zhang, L. (2019). Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance . In *Frontiers in Plant Science* (Vol. 10, p. 1068). <https://www.frontiersin.org/article/10.3389/fpls.2019.01068>
- Boutasknit, A., Baslam, M., Ait-El-Mokhtar, M., Anli, M., Ben-Laouane, R., Douira, A., El Modafar, C., Mitsui, T., Wahbi, S., & Meddich, A. (2020). Arbuscular Mycorrhizal Fungi Mediate Drought Tolerance and Recovery in Two Contrasting Carob (*Ceratonia siliqua* L.) Ecotypes by Regulating Stomatal, Water Relations, and (In)Organic Adjustments. *Plants*, 9(1). <https://doi.org/10.3390/plants9010080>
- Chun, S. C., Paramasivan, M., & Chandrasekaran, M. (2018). Proline accumulation influenced by osmotic stress in arbuscular mycorrhizal symbiotic plants. *Frontiers in Microbiology*, 9, 2525.
- Daryanti, D., Haryuni, H., & Utami, D. S. (2014). Pengimbasan Ketahanan Rhizoctonia Binukleat Terhadap Cekaman air Pada Bibit Vanili (*Vanilla Planifolia* Andrews). *Jurnal*

**Commented [A36]:** • References contain at least 20-25 references (80%) refers to the primary references / research journals and the latest sources (at least at last 10 years)

*Ilmiah Agrineca*, 14(1).

- Fileccia, T., Hovhera, V. ., & Bernoux, M. (2014). *Ukraine: Soil fertility to strengthen climate resilience preliminary assessment of the potential benefits of conservation agriculture*. Food and Agriculture Organization of the United Nations.
- Furlan, A. L., Bianucci, E., Giordano, W., Castro, S., & Becker, D. F. (2020). Proline metabolic dynamics and implications in drought tolerance of peanut plants. *Plant Physiology and Biochemistry*, 151, 566–578. <https://doi.org/https://doi.org/10.1016/j.plaphy.2020.04.010>
- Handajaningih, M., Marwanto, T., Armitasari, E., Inorih., & Pujiwati, H. (2018). Growth and yield response of melon to different time application of goat manure and dose of potassium. *Akta Agrosia*, 21(1), 1–5.
- Harahap, F. ., & Hidayat, B. (2021). The effect of rice husk ash and goat manure on the growth of spinach (*Amaranthus gangetius*). *Jurnal Pertanian Tropik*, 8(1), 26–32.
- Haryuni. (2015 a). The Effect of Hot Water Treatment and Dose Trichoderma sp. to Plant Tissue of Seedling Growth from Bud Chips of Sugarcane (*Saccharum officinarum*). *Biosaintifika Journal of Biology & Biology Education*, 7(1), 1–7. <https://doi.org/10.15294/biosaintifika.v7i1.3541>
- Haryuni, H., Dewi, T. S. K., & Nuryati, T. (2015 b). Pengaruh Dosis Rhizoctonia Binukleat (BNR) dan Pupuk posfor terhadap Pertumbuhan benih Vanili (*Vanilla planifolia* Andrew). *Prosiding Seminar Nasional & Internasional*. 36-45.
- Haryuni, H., Harahap, A. F. P., Supartini, Priyatmojo, A., & Gozan, M. (2020). The Effects of Biopesticide and *Fusarium oxysporum* f.sp. *vanillae* on the Nutrient Content of Binucleate *Rhizoctonia* -Induced Vanilla Plant . *International Journal of Agronomy*, 2020, 1–6. <https://doi.org/10.1155/2020/5092893>
- Haryuni, H., Suprpti, E., Dewi, T. S. K., Supriyadi, T., Nugroho, A. A., Priyatmojo, A., & Gozan, M. (2018). Phosphorus dosage and cow urine to chlorophyll and proline content on binucleate *Rhizoctonia* by induced resistance of vanilla. *Proceedings of the International Conference on Science and Education and Technology*, 247, 215–218.
- Havkin-Frenkel, D., & Belanger, F. . (2018). *Handbook Of Vanilla Science And Technology*. *Food Science And Technology 2nd*, Wiley Blackwell.
- Hendrati, R. L., Rachmawati, D., & Pamuji, A. C. (2016). Drought responses on growth, proline content and root anatomy of *Acacia auriculiformis* Cunn., *Tectona grandis* L., *Alstonia spectabilis* Br., and *Cedrela odorata* L. *Jurnal Penelitian Kehutanan Wallacea*, 5(2), 123–133.

- Huang, W., Ratkowsky, D. A., Hui, C., Wang, P., Su, J., & Shi, P. (2019). Leaf Fresh Weight Versus Dry Weight: Which is Better for Describing the Scaling Relationship between Leaf Biomass and Leaf Area for Broad-Leaved Plants? *Forests*, *10*(3).  
<https://doi.org/10.3390/f10030256>
- Hughes, A. P., Cockshull, K. E., & Heath, O. V. S. (1970). Leaf Area and Absolute Leaf Water Content. *Annals of Botany*, *34*(2), 259–266.  
<https://doi.org/10.1093/oxfordjournals.aob.a084366>
- Ichsan, M. C., Santoso, I., & Oktarina, O. (2016). Uji Efektivitas Waktu Aplikasi Bahan Organik dan Dosis Pupuk Sp-36 dalam Meningkatkan Produksi Okra (*Abelmoschus Esculentus*). *Agritrop: Jurnal Ilmu-Ilmu Pertanian (Journal of Agricultural Science)*, *14*(2).
- Lin, S., Shao, L., Hui, C., Song, Y., Reddy, G. V. P., Gielis, J., Li, F., Ding, Y., Wei, Q., & Shi, P. (2018). Why Does Not the Leaf Weight-Area Allometry of Bamboos Follow the 3/2-Power Law? *Frontiers in Plant Science*, *9*. <https://doi.org/10.3389/fpls.2018.00583>
- Muslim, A. (2019). *Pengendalian Hayati Patogen Tanaman Dengan Mikroorganisme Antagonis*. Unsri Press. <https://repository.unsri.ac.id/8930/>
- Nurcholis. (2017). Perbanyakkan tanaman panili (*Vanilla planifolia* Andrews) secara stek dan upaya untuk mendukung keberhasilan serta pertumbuhannya. *Agrovigor*, *10*(2), 149–156.
- Osakabe, Y., Osakabe, K., Shinozaki, K., & Tran, L.-S. P. (2014). Response of plants to water stress. *Frontiers in Plant Science*, *5*, 86.
- Pardosi, D. (2021). *Penampilan Morfologi Beberapa Varietas Tanaman Cabai Rawit (Capsicum frutescens L.) terhadap Cekaman Kekeringan*.
- Pratomo, H., & Prasetyo, B. (2018). Pembuatan pupuk kompos berbahan feses kambing menggunakan bantuan effective microorganism (EM4), Di Desa Tegal, Bogor. *Prosiding PKM-CSR*, *1*(2018), 403–412.
- Prihastati, E. (2012). Peranan dan Pola Akumulasi Proline Tanaman pada Adaptasi Cekaman Kekeringan. *Prosiding Seminar Biologi*, *9*(1), 594–597.
- Rasmussen, H. N. (1995). Orchid mycorrhiza. In *Terrestrial Orchids* (pp. 142–173). Cambridge University Press. <https://doi.org/10.1017/CBO9780511525452.009>
- Singh, R. P. (2012). Organic fertilizers: Types, production and environmental impact. In *Organic Fertilizers: Types, Production and Environmental Impact* (Issue July 2012).
- Siregar, B. L. (2015). Prolin: Metabolisme Dan Peranannya Dalam Tanaman Stress Kekeringan. *Jurnal Penelitian Ilmu Pertanian Methadagro*, *1*(1), 1–8.  
[https://uhn.ac.id/content.php?ref=TXgL&ssid=3\\_210204112326](https://uhn.ac.id/content.php?ref=TXgL&ssid=3_210204112326)

- Subantoro, R. (2014). Pengaruh cekaman kekeringan terhadap respon fisiologis perkecambahan benih kacang tanah (*Arachis hypogaea* L). *Mediagro*, 10(2).
- Surdianto, Y., Sutrisna, N., Basuno., & Solihi. (2015). Panduan teknis cara membuat arang sekam padi. *Balai Pengkajian Teknologi Pertanian (BPTP) Jawa Barat*, 23.  
<http://repository.pertanian.go.id/handle/123456789/6751>
- Tapwal, A., Kumar, R., & Borah, D. (2015). Effect of mycorrhizal inoculations on the growth of *Shorea robusta* seedlings. *Nusantara Bioscien*, 7(1), 1–5.
- Widiastuti, E., & Latifah, E. (2016). Keragaan Pertumbuhan dan Biomassa Varietas Kedelai (*Glycine max* (L)) di Lahan Sawah dengan Aplikasi Pupuk Organik Cair. *Jurnal Ilmu Pertanian Indonesia (JIPI)*, 21(2), 90–97.
- Yenni, M., Ibrahim, M. H., Nulit, R., & Sakimin, S. Z. (2021). The interactive effects of fertilizer and water stress on plant growth, leaf gas exchange and nutrient uptake on strawberry (*Fragaria x ananassa*, Duch). *AIMS Environmental Science*, 8(6), 597–618.
- Zhang, S., Wang, L., Ma, F., Zhang, X., & Fu, D. (2016). Arbuscular mycorrhiza improved phosphorus efficiency in paddy fields. *Ecological Engineering*, 95, 64–72.  
<https://doi.org/https://doi.org/10.1016/j.ecoleng.2016.06.029>



HOME ABOUT USER HOME SEARCH CURRENT ARCHIVES ANNOUNCEMENTS CONFERENCE PUBLICATION COLLABORATION

Home > User > Author > Submissions > #35618 > **Editing**

## #35618 Editing

SUMMARY REVIEW **EDITING**

### Submission

Authors Haryuni Haryuni, Muhamad Amin, Endang Suprapti, Tyas Soemarah Kurnia Dewi, Eko Hartoyo

Title Proline Reduction and Increasing Growth of Vanilla Plants Induced by BNR Fungus with Dose of Goat Manure and Husk Charcoal

Section Articles

Editor Dyah Indriyanti

### Copyediting

COPYEDIT INSTRUCTIONS

REVIEW METADATA	REQUEST	UNDERWAY	COMPLETE
1. Initial Copyedit File: None	—	—	—
2. Author Copyedit File: None <input type="button" value="Choose File"/> No file chosen <input type="button" value="Upload"/>	—	—	
3. Final Copyedit File: None	—	—	—

Copyedit Comments No Comments

### Layout

Galley Format	FILE
1. PDF VIEW PROOF	35618-91540-2-PB.PDF 2022-04-06 276

Supplementary Files	FILE
1. PROLINE REDUCTION AND INCREASING GROWTH OF VANILLA PLANTS INDUCED BY BNR FUNGUS WITH DOSE OF GOAT MANURE AND HUSK CHARCOAL	35618-90510-2-SP.DOCX 2022-03-14

Layout Comments No Comments

### Proofreading

REVIEW METADATA	REQUEST	UNDERWAY	COMPLETE
1. Author	—	—	
2. Proofreader	—	—	—
3. Layout Editor	—	—	—

Proofreading Corrections No Comments [PROOFING INSTRUCTIONS](#)




This work is licensed under a Creative Commons Attribution 4.0 International License.



#### ABOUT THE JOURNAL

- [Aims and Scope](#)
- [Publication Ethics](#)
- [Indexing & Abstracting](#)
- [Editorial Team](#)
- [Contact](#)

#### COLLABORATE WITH



#### FOR AUTHORS

- [Guidelines for Author](#)
- [Manuscript Template](#)
- [Mendeley Guidelines](#)
- [Scopus Citation Analysis](#)

#### Readers

ID 328,827	PK 1,146
US 21,887	VN 1,059
CN 11,072	IR 1,008
PH 10,460	NG 1,007
IN 6,316	GB 975
MY 4,142	MX 825
SG 2,693	AU 802
TH 2,256	KR 796
BR 1,606	CA 740
JP 1,406	EG 737
TR 1,169	TW 656

Pageviews: 1,077,522  
Flags Collected: 198

[View My Stats](#)

#### USER

You are logged in as...

**haryuni**

- [» My Journals](#)
- [» My Profile](#)
- [» Log Out](#)





## INVOICE

<https://journal.unnes.ac.id/nju/index.php/biosaintifika>  
[biosaintifika@mail.unnes.ac.id](mailto:biosaintifika@mail.unnes.ac.id)

Invoice : IV-1-12-2022  
Invoice Date : March 29, 2022

Amount due:  
1,500,000 (IDR)

Bill to:

**Haryuni Haryuni, Muhammad Amin, Endang Suprapti, Tyas Sumarah Kurnia Dewi, Eko Hartoyo**

Description	Amount
<b>Publication fee:</b> <b>Proline Reduction and Increasing Growth of Vanilla Plants Induced By BNR Fungus With Dose of Goat Manure and Husk Charcoal</b>	
Translate fee	1,500,000 (IDR) - 0 (IDR)
	<b>Subtotal</b> 1,500,000 (IDR)
	<b>Total</b> 1,500,000 IDR

Please send your payment to **BNI, No. 8581010420220013, Haryuni**


### Notes

Dear

Haryuni Haryuni, Muhammad Amin, Endang Suprapti, Tyas Sumarah Kurnia Dewi, Eko Hartoyo

Based on the results of the first review conducted by editorial team of Biosaintifika, your manuscript is ACCEPTED and it will be continued to our external reviewer. The publication fee for your accepted paper for volume 14 Issue 1, April 2022 is due on April 7, 2022

Biosaintifika Indexed by:





## Transaksi Berhasil

Nomor Rekening Tujuan \*\*\*\*\*013

Tanggal Transaksi 30-03-2022

Waktu Transaksi 11:49:41 WIB

Nama Pelanggan Haryuni

Nomor Rekening Pengirim \*\*\*\*\*453

Nomor Jurnal 915898

Nominal Rp1.500.000,00

Fee Rp0,00

Total Rp1.500.000,00

Keterangan UNIVERSITAS NEGERI SEMARANG